

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/00, 15/09, C07H 21/02, 21/04	A1	(11) International Publication Number: WO 98/40477 (43) International Publication Date: 17 September 1998 (17.09.98)
(21) International Application Number: PCT/US98/04978 (22) International Filing Date: 13 March 1998 (13.03.98) (30) Priority Data: 08/818,252 14 March 1997 (14.03.97) US 08/818,253 14 March 1997 (14.03.97) US 08/919,143 27 August 1997 (27.08.97) US (71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612-3550 (US). (72) Inventors: TSIEN, Roger, Y.; 8535 Nottingham Place, La Jolla, CA 92037 (US). MIYAWAKI, Atsushi; 3899 Nobel Drive, San Diego, CA 92122 (US). (74) Agent: HAILE, Lisa, A.; Fish & Richardson P.C., Suite 1400, 4225 Executive Square, La Jolla, CA 92037 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: FLUORESCENT PROTEIN SENSORS FOR DETECTION OF ANALYTES (57) Abstract Fluorescent indicators including a binding protein moiety, a donor fluorescent protein moiety, and an acceptor fluorescent protein moiety are described. The binding protein moiety has an analyte-binding region which binds an analyte and causes the indicator to change conformation upon exposure to the analyte. The donor moiety and the acceptor moiety change position relative to each other when the analyte binds to the analyte-binding region. The donor moiety and the acceptor moiety exhibit fluorescence resonance energy transfer when the donor moiety is excited and the distance between the donor moiety and the acceptor moiety is small. The indicators can be used to measure analyte concentrations in samples, such as calcium ion concentrations in cells.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

WO 98/40477

PCT/US98/04978

- 1 -

FLUORESCENT PROTEIN SENSORS FOR DETECTION OF ANALYTES

Statement as to Federally Sponsored Research

- 5 This invention was made with Government support under Grant No. NS27177, awarded by the National Institutes of Health. The Government has certain rights in this invention.

Background of the Invention

The invention relates to fluorescent protein sensors for detecting and quantifying analytes.

- 10 Measurement of an analyte concentration *in vitro* or *in vivo* by non-invasive techniques can help elucidate the physiological function of the analyte. This can also aid in identifying changes that occur in a cell or organism in response to physiological stimuli. For example, cyclic AMP can be detected by fluorescence resonance energy transfer between a separately labeled proteins that associate with each other but are not
15 covalently attached to each other. See, U.S. Pat. No. 5,439,797.

For example, many effects of Ca^{2+} in cells are mediated by Ca^{2+} binding to calmodulin (CaM), which causes CaM to bind and activate target proteins or peptide sequences. Based on the NMR solution structure of CaM bound to the 26-residue M13 Ca^{2+} -binding peptide of myosin light-chain kinase, Porumb *et al.* fused the C-terminus of CaM via a
20 Gly-Gly spacer to the M13. Ca^{2+} binding switches the resulting hybrid protein (CaM-M13) from a dumbbell-like extended form to a compact globular form similar to the CaM-M13 intermolecular complex. See, Porumb, T., *et al.*, *Prot.Engineering* 7:109-115 (1994).

Fluorescent Ca^{2+} indicators such as fura-2, indo-1, fluo-3, and Calcium-Green have been
25 the mainstay of intracellular Ca^{2+} measurement and imaging. See, for example, U.S. Pat. No. 4,603,209 and U.S. Pat. No. 5,049,673. These relatively low molecular weight indicators can suffer from many technical problems relating to ester loading, leakage of

- 2 -

5

This invention provides fluorescent indicators and methods for using them to determine the concentration of an analyte both *in vitro* and *in vivo*. In one aspect, the fluorescent indicator includes a binding protein moiety, a donor fluorescent protein moiety, and an acceptor fluorescent protein moiety. The binding protein moiety has an analyte-binding region which binds an analyte and causes the indicator to change conformation upon exposure to the analyte. The donor fluorescent protein moiety is covalently coupled to the binding protein moiety. The acceptor fluorescent protein moiety is covalently coupled to the binding protein moiety. In the fluorescent indicator, the donor moiety and the acceptor moiety change position relative to each other when the analyte binds to the analyte-binding region, altering fluorescence resonance energy transfer between the donor moiety and the acceptor moiety when the donor moiety is excited.

The donor fluorescent protein moiety and the acceptor fluorescent protein moiety can be *Aequorea*-related fluorescent protein moieties. Preferably, the donor fluorescent protein moiety is P4-3, EBFP, or W1B, and the acceptor fluorescent protein moiety is S65T, EGFP, or 10c.

In preferred embodiments, the indicator further includes the target peptide moiety and a linker moiety that covalently couples the binding protein and the target peptide moiety. The binding protein moiety further includes a peptide-binding region for binding the target peptide moiety. The binding protein moiety can be covalently coupled to the donor fluorescent protein moiety and the target peptide moiety can be covalently coupled to the acceptor fluorescent protein moiety.

WO 98/40477

PCT/US98/04978

- 3 -

The indicator can be a single polypeptide. In preferred embodiments, one of the donor fluorescent protein moiety or the acceptor fluorescent protein moiety is covalently coupled to the carboxy terminus of the single polypeptide and the other of the donor fluorescent protein moiety or the acceptor fluorescent protein moiety is covalently
5 coupled to the amino terminus of the single polypeptide.

The indicator can include a localization sequence. The localization sequence can be a nuclear localization sequence, an endoplasmic reticulum localization sequence, a peroxisome localization sequence, a mitochondrial import sequence, a mitochondrial localization sequence, or a localized protein.

- 10 In preferred embodiments, the linker moiety is a peptide moiety. The linker moiety can include between about 1 amino acid residue and about 20 amino acid residues. The linker moiety can be -Gly-Gly-.

Preferably, the binding protein moiety is calmodulin, a calmodulin-related protein moiety, cGMP-dependent protein kinase, a steroid hormone receptor, a ligand binding
15 domain of a steroid hormone receptor, protein kinase C, inositol-1,4,5-triphosphate receptor, or recoverin. A calmodulin-related protein moiety is derived from calmodulin that has been modified to have a different binding affinity for calcium or a target peptide moiety.

Most preferably, the binding protein moiety is calmodulin or a calmodulin-related protein
20 moiety. In these embodiments, the target peptide moiety can be a subsequence of a calmodulin-binding domain of M13, smMLCKp, CaMKII, Caldesmon, Calspermin, Calcineurin, PhK5, PhK13, C28W, 59-kDa PDE, 60-kDa PDE, NO-30, AC-28, *Bordetella pertussis* AC, Neuromodulin, Spectrin, MARCKS, F52, β -Adducin, HSP90a, HIV-1 gp160, BBMHBI, Dilute MHC, Mastoparan, Melittin, Glucagon, Secretin, VIP,
25 GIP, or Model Peptide CBP2. Preferably, the target peptide moiety is M13.

WO 98/40477

PCT/US98/04978

- 4 -

In another aspect, the invention features a fluorescent indicator including a target peptide moiety, a binding protein moiety, a linker moiety, a donor fluorescent protein moiety covalently coupled to the binding protein moiety, and an acceptor fluorescent protein moiety covalently coupled to the binding protein moiety. The binding protein moiety has
5 an analyte-binding region which binds an analyte and causes the indicator to change conformation upon exposure to the analyte. The linker moiety covalently couples the binding protein and the target peptide moiety and is a peptide moiety. The binding protein moiety has a peptide-binding region for binding the target peptide moiety. The indicator is a single polypeptide.

10 In another aspect, the invention features a method for determining the concentration of an analyte in a sample. The method includes the steps of contacting the sample with a fluorescent indicator having a donor fluorescent protein moiety, binding protein moiety, and acceptor protein moiety, exciting the donor moiety, and determining the degree of
15 fluorescence resonance energy transfer in the sample corresponding to the concentration of the analyte in the sample.

In preferred embodiments, the step of determining the degree of fluorescence resonance energy transfer in the sample includes measuring light emitted by the acceptor fluorescent protein moiety. In other preferred embodiments, determining the degree of fluorescence resonance energy transfer in the sample includes measuring light emitted
20 from the donor fluorescent protein moiety, measuring light emitted from the acceptor fluorescent protein moiety, and calculating a ratio of the light emitted from the donor fluorescent protein moiety and the light emitted from the acceptor fluorescent protein moiety. In yet other preferred embodiments, determining the degree of fluorescence resonance energy transfer in the sample includes measuring the excited state lifetime of
25 the donor moiety.

The method can further include the steps of determining the concentration of the analyte at a first time after contacting the sample with the fluorescence indicator, determining the concentration of the analyte at a second time after contacting the sample with the

WO 98/40477

PCT/US98/04978

- 5 -

fluorescence indicator, and calculating the difference in the concentration of the analyte at the first time and the second time, whereby the difference in the concentration of the analyte in the sample reflects a change in concentration of the analyte present in the sample.

- 5 In other embodiments, the method can further include the step of contacting the sample with a compound between the first time and the second time, whereby a difference in the concentration of the analyte in the sample between the first time and the second time indicates that the compound alters the presence of the analyte.

In preferred embodiments, the sample includes an intact cell and the contacting step
10 includes incorporating the fluorescent indicator into the cell. The step of incorporating the fluorescent indicator into the cell can include transfecting the cell with an expression vector comprising expression control sequences operably linked to a nucleic acid sequence coding for the expression of the fluorescent indicator. The preferred analyte is calcium.

- 15 In yet another aspect, the invention features an isolated nucleic acid sequence which encodes the fluorescent indicator. In preferred embodiments, an expression vector or a transgenic non-human animal includes the nucleic acid sequence.

In another aspect, the invention features an expression vector including expression control sequences operatively linked to a nucleic acid sequence coding for the expression
20 of the fluorescent indicator. The expression vector can be adapted for function in a prokaryotic cell or a eukaryotic cell.

In another aspect of the invention, a host cell transfected with an expression vector can include an expression control sequence operatively linked to a sequence coding for the expression of the fluorescent indicator. The cell can be a prokaryote, such as *E. coli*, or
25 a eukaryotic cell, such as a yeast cell or mammalian cell.

WO 98/40477

PCT/US98/04978

- 6 -

In another aspect, the invention features a transgenic non-human animal having a phenotype characterized by expression of the nucleic acid sequence coding for the expression of the fluorescent indicator. The phenotype is conferred by a transgene contained in the somatic and germ cells of the animal. The animal can be a mouse.

- 5 In another aspect, the invention features a method for producing a transgenic non-human animal having a phenotype characterized by expression of the nucleic acid sequence coding for the expression of the fluorescent indicator. The method includes the steps of: (a) introducing a transgene into a zygote of an animal, the transgene comprising a DNA construct encoding the fluorescent indicator; (b) transplanting the zygote into a
- 10 pseudopregnant animal; (c) allowing the zygote to develop to term; and (d) identifying at least one transgenic offspring containing the transgene. The step of introducing of the transgene into the embryo can be by introducing an embryonic stem cell containing the transgene into the embryo, or infecting the embryo with a retrovirus containing the transgene.
- 15 "Peptide" refers to a polymer in which the monomers are amino acid residues which are joined together through amide bonds, alternatively referred to as a polypeptide. A "single polypeptide" is a continuous peptide that constitutes the protein. When the amino acids are alpha-amino acids, either the L-optical isomer or the D-optical isomer can be used, the L-isomers being preferred. Additionally, unnatural amino acids such as beta-
- 20 alanine, phenylglycine, and homoarginine are meant to be included. Commonly encountered amino acids which are not gene-encoded can also be used in the present invention, although preferred amino acids are those that are encodable. For a general review, see, for example, Spatola, A.F., in *Chemistry and Biochemistry of Amino Acids, Peptides and Proteins*, B. Weinstein, ed., Marcel Dekker, New York, p. 267 (1983).
- 25 "Fluorescent protein" refers to any protein capable of emitting light when excited with appropriate electromagnetic radiation. Fluorescent proteins include proteins having amino acid sequences that are either natural or engineered, such as the fluorescent proteins derived from *Aequorea*-related fluorescent proteins.

WO 98/40477

PCT/US98/04978

- 7 -

In FRET, the "donor fluorescent protein moiety" and the "acceptor fluorescent protein moiety" are selected so that the donor and acceptor moieties exhibit fluorescence resonance energy transfer when the donor moiety is excited. One factor to be considered in choosing the donor/acceptor fluorescent protein moiety pair is the efficiency of FRET
5 between the two moieties. Preferably, the efficiency of FRET between the donor and acceptor moieties is at least 10%, more preferably at least 50%, and most preferably at least 80%. The efficiency of FRET can be tested empirically using the methods described herein and known in the art, particularly, using the conditions set forth in the Examples.

10 "Covalently coupled" refers to a covalent bond or other covalent linkage between two moieties. The covalent linkage can include a diradical moiety linking to two moieties.

"Binding protein" refers to a protein capable of binding an analyte. Preferred binding proteins change conformation upon binding the analyte. "Target peptide" refers to a peptide that can bind to the binding protein. The target peptide can be a subsequence of
15 a peptide that binds to the binding protein.

"Analyte" refers to a molecule or ion in solution that binds to the binding protein, causing it to change conformation. Preferably, the analyte binds reversibly to the binding protein.

"Moiety" refers to a radical of a molecule that is attached to another radical of the indicator. Thus, a "fluorescent protein moiety" is the radical of a fluorescent protein
20 coupled to a binding protein moiety or a linker moiety, a "binding protein moiety" is a radical of a binding protein coupled to a fluorescent protein moiety, a "target peptide moiety" is a radical of a target peptide of the binding protein, and a "linker moiety" refers to the radical of a molecular linker that is ultimately coupled to both the donor and acceptor fluorescent protein moieties.

25 "Operatively linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. A control

WO 98/40477

PCT/US98/04978

- 8 -

sequence operatively linked to a coding sequence is ligated such that expression of the coding sequence is achieved under conditions compatible with the control sequences. "Control sequence" refers to polynucleotide sequences which are necessary to effect the expression of coding and non-coding sequences to which they are ligated. Control
5 sequences generally include promoter, ribosomal binding site, and transcription termination sequence. The term "control sequences" is intended to include, at a minimum, components whose presence can influence expression, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences.

- 10 "Polynucleotide" refers to a polymeric form of nucleotides of at least 10 bases in length. The nucleotides can be ribonucleotides, deoxynucleotides, or modified forms of either type of nucleotide. The term includes single and double stranded forms of DNA.

The invention can have one or more of the following advantages. Ligand-induced conformational changes can be monitored by FRET if, for example, the amino and
15 carboxy termini of the binding protein are fused to a donor and acceptor GFP. This approach has several advantages over the usual covalent labeling with fluorescent probes. The fluorescent indicator can be generated *in situ* by gene transfer into the cells or organisms. This approach avoids the need to express and purify large quantities of soluble recombinant protein, purify and label it *in vitro*, microinject it back into cells.
20 The fluorescent indicator can be targeted to cellular structures. The sites of fusion between the moieties of the fluorescent indicator are exactly defined, giving a molecularly homogenous product without relying on elaborate protein chemistry. In addition, the chromophore of GFP is fixed in the protein. See, Ormo, M., *et al.*, *Science* 273:1392-1395 (1996). If the GFP donor and acceptor are fused to a host protein rigidly,
25 minor changes in the relative orientation of the ends of the latter would alter FRET. In contrast, most conventional fluorescent labels are attached by flexible linkers that at least partially decouple the fluorophore orientation from

WO 98/40477

PCT/US98/04978

- 9 -

that of the protein to which it is attached, limiting the sensitivity of the FRET measurement.

Other features or advantages of the present invention will be apparent from the following detailed description of the invention, and also from the claims.

5 Brief Description of the Drawings

FIG. 1 is a schematic diagram depicting a fluorescent indicator that measures the concentration of an analyte by fluorescence resonance energy transfer.

FIG. 2a is a schematic diagram depicting the structure of chimera proteins containing P4-3, CaM-M13, and S65T, where HIS is the amino-terminal tag peptide containing the
10 polyhistidine sequence and XCaM is Xenopus calmodulin.

FIG. 2b is a diagram depicting the amino acid and nucleotide sequences of the boundaries between P4-3 and CaM and between M13 and S65T in cameleon-1.

FIG. 3 is a graph depicting the emission spectra of cameleon-1 before and after addition of 2 mM CaCl_2 to give 1 mM free Ca^{2+} .

15 FIG. 4a is a graph depicting the change in emission spectrum of cameleon-1 on titration with Ca^{2+} when excited at 380 nm.

FIG. 4b is a graph depicting Ca^{2+} titration curves of cameleon-1 (open circles) and cameleon-1/E104Q (solid circles).

FIG. 5 is a schematic diagram depicting the structures of cameleon-2 and cameleon-3,
20 and their derivatives, cameleon-2nu and cameleon-3er, where Kz is Kozak's consensus sequence.

WO 98/40477

PCT/US98/04978

- 10 -

FIG. 6a is a digital fluorescence image depicting fluorescence of cameleon-2 localized to the cytosol of HeLa cells, where the bar is 10 Tm.

FIG. 6b is a graph depicting temporal changes in the emission ratio of cameleon-2 for each of the HeLa cells shown in FIG. 6a.

- 5 FIG. 6c is a digital fluorescence image depicting fluorescence of cameleon-2nu localized to the nuclei of HeLa cells, where the bar is 10 Tm.

FIG. 6d is a graph depicting temporal changes in the emission ratio of cameleon-2nu in the two nuclei shown in FIG. 6c.

- FIG. 6e is a digital fluorescence image depicting fluorescence of cameleon-3er in
10 transfected HeLa cells, where the bar is 10 Tm.

FIG. 6f is a graph depicting the time-course of emission ratio of cameleon-3er (average of four cells).

FIG. 7 depicts the nucleotide sequence (SEQ ID NO:1) and amino acid sequence (SEQ ID NO:2) of cameleon-2.

- 15 FIG. 8 depicts the nucleotide sequence (SEQ ID NO:3) and amino acid sequence (SEQ ID NO:4) of cameleon-2nu.

FIG. 9 depicts the nucleotide sequence (SEQ ID NO:5) and amino acid sequence (SEQ ID NO:6) of cameleon-3.

- FIG. 10 is a list depicting the nucleotide sequence (SEQ ID NO:7) and amino acid
20 sequence of cameleon-3er (SEQ ID NO:8).

WO 98/40477

PCT/US98/04978

- 11 -

Detailed Description

A fluorescent indicator that utilizes fluorescent resonance energy transfer ("FRET") to measure the concentration of an analyte includes two fluorescent protein moieties having emission and excitation spectra that render one a donor fluorescent protein moiety and the other an acceptor fluorescent protein moiety. The fluorescent protein moieties are chosen such that the excitation spectrum of one of the moieties (the acceptor fluorescent protein moiety) overlaps with the emission spectrum of the excited protein moiety (the donor fluorescent protein moiety). The donor and acceptor fluorescent protein moieties are bound to a binding protein moiety that changes conformation upon binding the analyte. The change in conformation leads to a change in relative position and orientation of the donor and acceptor fluorescent protein moieties, thereby altering the relative amounts of fluorescence from the two fluorescent protein moieties when the donor is excited by irradiation. In particular, binding of the analyte changes the ratio of the amount of light emitted by the donor and acceptor fluorescent protein moieties. The ratio between the two emission wavelengths provides a measure of the concentration of the analyte in the sample, which is based in part on the binding affinity of the binding protein moiety and the analyte.

Referring to FIG. 1, the donor fluorescent protein moiety is covalently linked to a first region (e.g., the amino terminus) of the binding protein moiety, and the acceptor fluorescent protein moiety is covalently linked to a second region (e.g., the carboxy terminus) of the binding protein moiety such that the donor and acceptor moieties move closer together upon binding the analyte. Alternatively, the donor and acceptor moieties can move farther apart upon binding the analyte. In one embodiment, depicted in FIG. 1, the acceptor moiety is covalently bonded to a target peptide moiety that also binds to the binding protein moiety and the target peptide moiety is covalently bonded to the binding protein moiety by a linker moiety. The linker moiety is flexible enough to allow the target peptide moiety to bind to the binding protein moiety. The donor moiety is excited by light of appropriate intensity within the excitation spectrum of the donor moiety ($\Sigma_{\text{excitation}}$). The donor moiety emits the absorbed energy as fluorescent light (Σ_{emission}). When the acceptor fluorescent protein moiety is positioned to quench the

WO 98/40477

PCT/US98/04978

- 12 -

donor moiety in the excited state, the fluorescence energy is transferred to the acceptor moiety which can emit fluorescent light ($\Sigma_{\text{emission } 2}$). FRET can be manifested as a reduction in the intensity of the fluorescent signal from the donor moiety ($\Sigma_{\text{emission } 1}$), reduction in the lifetime of the excited state of the donor moiety, or emission of
5 fluorescent light at the longer wavelengths (lower energies) characteristic of the acceptor moiety ($\Sigma_{\text{emission } 2}$). When the conformation of the binding protein moiety changes upon binding the analyte, the fluorescent protein moieties come closer together (or physically separate), and FRET is increased (or decreased) accordingly.

The efficiency of FRET depends on the separation distance and the orientation of the
10 donor and acceptor fluorescent protein moieties. For example, the Forster equation describes the efficiency of excited state energy transfer, based in part on the fluorescence quantum yield of the donor moiety and the energetic overlap with the acceptor moiety. The Forster equation is:

$$E = (F_0 - F)/F_0 = R_0^6 / (R^6 + R_0^6)$$

15 where E is the efficiency of FRET, F and F_0 are the fluorescence intensities of the donor moiety in the presence and absence of the acceptor, respectively, and R is the distance between the donor moiety and the acceptor moiety.

The characteristic distance R_0 at which FRET is 50% efficient depends on the quantum yield of the donor moiety (*i.e.*, the shorter-wavelength fluorophore), the extinction
20 coefficient of the acceptor moiety (*i.e.*, the longer-wavelength fluorophore), and the overlap between the emission spectrum of the donor moiety and the excitation spectrum of the acceptor moiety. R_0 is given (in D) by

$$R_0 = 9.79 \times 10^3 (K^2 Q J n^{-4})^{1/6}$$

where K^2 is an orientation factor having an average value close to 0.67 for freely mobile
25 donors and acceptors, Q is the quantum yield of the unquenched donor moiety, n is the refractive index of the medium separating the donor moiety and the acceptor moiety, and

WO 98/40477

PCT/US98/04978

- 13 -

J is the overlap integral. J can be quantitatively expressed as the degree of spectral overlap between the donor moiety and the acceptor moiety according to the equation:

$$J = I_0^4 M_{\Sigma} F_{\Sigma} \Sigma^4 d\Sigma / I_0^4 F_{\Sigma} d\Sigma$$

where M_{Σ} ($M^{-1}cm^{-1}$) is the molar absorptivity of the acceptor and F_{Σ} is the donor moiety
5 fluorescence intensity at wavelength Σ . See, for example, Forster, T. *Ann. Physik* 2:55-75
(1948). Tables of spectral overlap integrals are readily available to those working in the
field (for example, Berlman, I.B. *Energy transfer parameters of aromatic compounds*,
Academic Press, New York and London (1973)). FRET is a nondestructive
spectroscopic method that can monitor proximity and relative angular orientation of
10 fluorophores in living cells. See, for example, Adams, S.R., *et al.*, *Nature* 349:694-697
(1991), and Gonzalez, J. & Tsien, R.Y. *Biophys. J.* 69:1272-1280 (1995).

These factors need to be balanced to optimize the efficiency and detectability of FRET
from the fluorescent indicator. The emission spectrum of the donor fluorescent protein
moiety should overlap as much as possible with the excitation spectrum of the acceptor
15 fluorescent protein moiety to maximize the overlap integral J. Also, the quantum yield
of the donor fluorescent protein moiety and the extinction coefficient of the acceptor
fluorescent protein moiety should be as large as possible to maximize R_0 . In addition,
the excitation spectra of the donor and acceptor moieties should overlap as little as
possible so that a wavelength region can be found at which the donor moiety can be
20 excited selectively and efficiently without directly exciting the acceptor moiety. Direct
excitation of the acceptor moiety should be avoided since it can be difficult to distinguish
direct emission from fluorescence arising from FRET. Similarly, the emission spectra
of the donor and acceptor moieties should have minimal overlap so that the two
emissions can be distinguished. High fluorescence quantum yield of the acceptor moiety
25 is desirable if the emission from the acceptor moiety is to be monitored to determine
analyte concentration in a sample. In a preferred embodiment, the donor fluorescent
protein moiety is excited by ultraviolet (<400 nm) and emits blue light (<500 nm), and
the acceptor fluorescent protein moiety is efficiently excited by blue but not by

WO 98/40477

PCT/US98/04978

- 14 -

ultraviolet light and emits green light (>500 nm), for example, P4-3 and S65T, respectively.

In another preferred embodiment, the donor fluorescent moiety is excited by violet (400-430 nm) and emits blue-green (450-500 nm) and the acceptor fluorescent moiety is
5 efficiently excited by blue-green (450-500 nm) and emits yellow-green light (520 nm), for example WIB and 10C respectively.

The amount of analyte in a sample can be determined by determining the degree of FRET in the sample. Changes in analyte concentration can be determined by monitoring FRET at a first and second time after contact between the sample and the fluorescent indicator
10 and determining the difference in the degree of FRET. The amount of analyte in the sample can be calculated by using a calibration curve established by titration.

The degree of FRET can be determined by any spectral or fluorescence lifetime characteristic of the excited donor moiety. For example, intensity of the fluorescent signal from the donor, the intensity of fluorescent signal from the acceptor, the ratio of
15 the fluorescence amplitudes near the acceptor's emission maxima to the fluorescence amplitudes near the donor's emission maximum, or the excited state lifetime of the donor can be monitored.

Preferably, changes in the degree of FRET are determined as a function of the change in the ratio of the amount of fluorescence from the donor and acceptor moieties, a process
20 referred to as "ratioing." Changes in the absolute amount of indicator, excitation intensity, and turbidity or other background absorbances in the sample at the excitation wavelength affect the intensities of fluorescence from both the donor and acceptor approximately in parallel. Therefore the ratio of the two emission intensities is a more robust and preferred measure of cleavage than either intensity alone.

25 Fluorescence in a sample is measured using a fluorometer. In general, excitation radiation, from an excitation source having a first wavelength, passes through excitation

WO 98/40477

PCT/US98/04978

- 15 -

optics. The excitation optics cause the excitation radiation to excite the sample. In response, fluorescent proteins in the sample emit radiation which has a wavelength that is different from the excitation wavelength. Collection optics then collect the emission from the sample. The device can include a temperature controller to maintain the sample
5 at a specific temperature while it is being scanned. According to one embodiment, a multi-axis translation stage moves a microtiter plate holding a plurality of samples in order to position different wells to be exposed. The multi-axis translation stage, temperature controller, auto-focusing feature, and electronics associated with imaging and data collection can be managed by an appropriately programmed digital computer.
10 The computer also can transform the data collected during the assay into another format for presentation.

Methods of performing assays on fluorescent materials are well known in the art and are described in, e.g., Lakowicz, J.R., *Principles of Fluorescence Spectroscopy*, New York: Plenum Press (1983); Herman, B., Resonance energy transfer microscopy, in:
15 *Fluorescence Microscopy of Living Cells in Culture, Part B, Methods in Cell Biology*, vol. 30, ed. Taylor, D.L. & Wang, Y.-L., San Diego: Academic Press (1989), pp. 219-243; Turro, N.J., *Modern Molecular Photochemistry*, Menlo Park: Benjamin/Cummings Publishing Co., Inc. (1978), pp. 296-361.

The excited state lifetime of the donor moiety is, likewise, independent of the absolute
20 amount of substrate, excitation intensity, or turbidity or other background absorbances. Its measurement requires equipment with nanosecond time resolution.

Quantum yields of wild-type GFP, S65T, and P4-1 mutants can be estimated by comparison with fluorescein in 0.1 N NaOH as a standard of quantum yield 0.91. J.N. Miller, ed., *Standards in Fluorescence Spectrometry*, New York: Chapman and Hall
25 (1981). Mutants P4 and P4-3 were likewise compared to 9-aminoacridine in water (quantum yield 0.98).

WO 98/40477

PCT/US98/04978

- 16 -

Any fluorescent protein can be used in the invention, including proteins that fluoresce due intramolecular rearrangements or the addition of cofactors that promote fluorescence. For example, green fluorescent proteins of cnidarians, which act as their energy-transfer acceptors in bioluminescence, are suitable fluorescent proteins for use in the fluorescent indicators. A green fluorescent protein ("GFP") is a protein that emits green light, and a blue fluorescent protein ("BFP") is a protein that emits blue light. GFPs have been isolated from the Pacific Northwest jellyfish, *Aequorea victoria*, the sea pansy, *Renilla reniformis*, and *Phialidium gregarium*. See, Ward, W.W., *et al.*, *Photochem. Photobiol.*, 35:803-808 (1982); and Levine, L.D., *et al.*, *Comp. Biochem. Physiol.*, 72B:77-85 (1982).

A variety of *Aequorea*-related GFPs having useful excitation and emission spectra have been engineered by modifying the amino acid sequence of a naturally occurring GFP from *Aequorea victoria*. See, Prasher, D.C., *et al.*, *Gene*, 111:229-233 (1992); Heim, R., *et al.*, *Proc. Natl. Acad. Sci., USA*, 91:12501-04 (1994); U.S. Ser. No. 08/337,915, filed November 10, 1994; International application PCT/US95/14692, filed 11/10/95; and U.S. Ser. No. 08/706,408, filed August 30, 1996. The cDNA of GFP can be concatenated with those encoding many other proteins; the resulting fusions often are fluorescent and retain the biochemical features of the partner proteins. See, Cubitt, A.B., *et al.*, *Trends Biochem. Sci.* 20:448-455 (1995). Mutagenesis studies have produced GFP mutants with shifted wavelengths of excitation or emission. See, Heim, R. & Tsien, R.Y. *Current Biol.* 6:178-182 (1996). Suitable pairs, for example a blue-shifted GFP mutant P4-3 (Y66H/Y145F) and an improved green mutant S65T can respectively serve as a donor and an acceptor for fluorescence resonance energy transfer (FRET). See, Tsien, R.Y., *et al.*, *Trends Cell Biol.* 3:242-245 (1993). A fluorescent protein is an *Aequorea*-related fluorescent protein if any contiguous sequence of 150 amino acids of the fluorescent protein has at least 85% sequence identity with an amino acid sequence, either contiguous or non-contiguous, from the wild type *Aequorea* green fluorescent protein. More preferably, a fluorescent protein is an *Aequorea*-related fluorescent protein if any contiguous sequence of 200 amino acids of the fluorescent protein has at least 95% sequence identity with an amino acid sequence, either contiguous or non-contiguous, from the wild type *Aequorea* green fluorescent protein. Similarly, the fluorescent protein

WO 98/40477

PCT/US98/04978

- 17 -

can be related to *Renilla* or *Phialidium* wild-type fluorescent proteins using the same standards. Some *Aequorea*-related engineered versions described in Table I. Other variants or mutants are within the scope of the invention as described, for example, in the Examples.

5

TABLE I

<u>Clone</u>	<u>Mutation(s)</u>	<u>Excitation max</u> (nm)	<u>Emission</u> max (nm)	<u>Extinction Coefficient</u> (M ⁻¹ cm ⁻¹)	<u>Quantum</u> yield
<i>Wild type</i>	none	395 (475)	508	21,000 (7,150)	0.77
<i>P4</i>	Y66H	383	447	13,500	0.21
<i>P4-3</i>	Y66H;Y145F	381	445	14,000	0.38
10 <i>W7</i>	Y66W;N146I M153T V163A N212K	433 (453)	475 (501)	18,000 (17,100)	0.67
<i>W2</i>	Y66W;I123V Y145H H148R M153T V163A N212K	432 (453)	480	10,000 (9,600)	0.72
<i>S65T</i>	S65T	489	511	39,200	0.68
<i>P4-1</i>	S65T;M153A K238E	504 (396)	514	14,500 (8,600)	0.53
<i>S65A</i>	S65A	471	504		
15 <i>S65C</i>	S65C	479	507		
<i>S65L</i>	S65L	484	510		
<i>Y66F</i>	Y66F	360	442		
<i>Y66W</i>	Y66W	458	480		
<i>10c</i>	S65G;V68L S72A;T203Y	513	527		

WO 98/40477

PCT/US98/04978

- 18 -

<i>W1B</i>	F64L;S65T Y66W;N146I M153T V163A N212K	432 (453)	476 (503)
<i>Emerald</i>	S65T;S72A N149K M153T I167T	487	508

WO 98/40477

PCT/US98/04978

- 19 -

<i>Sapphire</i>	S72A;Y145F	395	511
	T203I		

An additional clone, W1B1 included the following mutations: F64L;S65T; Y66W; F99S; and V163A.

Other fluorescent proteins can be used in the fluorescent indicators, such as, for example, yellow fluorescent protein from *Vibrio fischeri* strain Y-1, Peridinin-chlorophyll *a* binding protein from the dinoflagellate *Symbiodinium* sp. phycobiliproteins from marine cyanobacteria such as *Synechococcus*, e.g., phycoerythrin and phycocyanin, or oat phytochromes from oat reconstructed with phycoerythrobilin. These fluorescent proteins have been described in Baldwin, T.O., *et al.*, *Biochemistry* 29:5509-5515 (1990), Morris, B.J., *et al.*, *Plant Molecular Biology*, 24:673-677 (1994), and Wilbanks, S.M., *et al.*, *J. Biol. Chem.* 268:1226-1235 (1993), and Li *et al.*, *Biochemistry* 34:7923-7930 (1995).

The efficiency of FRET between the donor and acceptor fluorescent protein moieties can be adjusted by changing ability of the two fluorescent proteins to closely associate. The nature of the binding protein moiety, target peptide moiety, and linker moiety each affect the FRET and the response of the indicator to the analyte. Generally, large conformational changes in the binding protein moiety are desired along with a high affinity for the target peptide moiety.

The binding protein moiety is a protein, or part thereof, that changes conformation upon binding an analyte. Proteins that undergo useful conformation change upon binding an analyte include calmodulin (CaM), cGMP-dependent protein kinase, steroid hormone receptors (or ligand binding domain thereof), protein kinase C, inositol-1,4,5-triphosphate receptor, or recoverin. See, for example, Katzenellenbogen, J.A. & Katzenellenbogen, B.S. *Chemistry & Biology* 3:529-536 (1996), and Ames, J.B., *et al.*, *Curr. Opin. Struct. Biol.* 6:432-438 (1996). The binding protein moiety preferably binds target peptides in addition to the analyte. The Ca²⁺-binding affinity of calmodulin can

WO 98/40477

PCT/US98/04978

- 20 -

be tuned as reviewed, for example, in Falke, J.J., *et al.*, *Quart. Rev. Biophys.* 27:219-290 (1994).

The target peptide moiety can contain any of the amino acid sequences in Table II, or a portion thereof with the proviso that the target peptide must bind to the binding protein moiety. The target peptide can be a subsequence of a calmodulin-binding domain. The target peptide moieties listed in Table II are recognized by the binding protein moiety CaM. See, for example, Crivici, A. & Ikura, M. *Annu. Rev. Biophys. Biomol. Struct.* 24:84-116 (1995). The target peptide moiety can be modified to enhance the response of the fluorescent indicator to the analyte. Other target peptide moieties are known in the art for other binding proteins.

WO 98/40477

PCT/US98/04978

- 21 -

TABLE II

Target ^a	Sequence
skMLCK (M13)	KRRWKKNFIAVSAANRFKKISSSGAL
smMLCK (smMLCKp)	ARRKWQKTGHAVRAIGRLSS
5 CaMKII	ARRKLKGAILTTMLATRNFS
Caldesmon	GVRNIKSMWEKGNVFSS
Calspermin	ARRKLKAAVKAVVASSRLGS
PFK (M11)	FMNNWEVYKLLAHIRPPAPKSGSYTV
Calcineurin	ARKEVIRNKIRAIGKMARVFSVLR
10 PhK (PhK5)	LRRLIDAYAFRIYGHVVKKGQQQNRG
(PhK13)	RGKFKVICLTVLASVRIYYQYRRVKPG
Ca ²⁺ -ATPase (C28W)	LRRGQILWFRGLNRIQTQIKVVNAFSSS
59-kDa PDE	RRKHLQRPIFRLRCLVKQLEK
60-kDa PDE	TEKMWQRLKGILRCLVKQLEK
15 NOS (NO-30)	KRRAIGFKKLAEAVKFSAKLMGQ
Type I AC (AC-28)	IKPAKRMKFKTVCYLLVQLMHCRKMFKA
<i>Borderella periussis</i> AC	IDLLWKIARAGARSAVGTEA
Neuromodulin	KAHKAATKIQASFRGHITRKKLKGEKK
Spectrin	KTASPWKSARLMVHTVATFNSIKE
20 MARCKS	KKKKKRFSFKKSFKLSGFSFKKSKK
F52 or MacMARKS	KKKKKFSSFKKPFKLSGLSFKRNRK
β-Adducin	KQQKEKTRWLNTPTNTYLRVNVADDEVQRNMGS
HSP90a	KDQVANSASFQERLRKHGLEVI
HIV-1 gp160	YHRLRDLLLIVKRIVELLGRR
25 BBMHBI	QQLATLIQKTYRGWRCRTHYQLM
Dilute MHC	RAACIRIQKTIRGWLLRKRYLCMQ
Mastoparan	INLKALAALAKKIL
Melittin	GIGAVLKVLTTGLPALISWIKRKRQQ
Glucagon	HSQGTFTTSDYSKYLDSSRAQDFVQWLMNT
30 Secretin	HSDGTFTSELSRLRDSARLQRLQLGLV
VIP	HSDAVFTDNYTRLRKQMAVKKYLNSILN

PCT/US98/04978

GIP YADGTFISDYS AIMNKIRQQDFVNWLLAQQQKS
Model Peptide CBP2 KLWKKLLKLLKLLKLG

The length of the linker moiety is chosen to optimize both FRET and the kinetics and specificity of conformational changes induced by analyte binding. The linker moiety should be long enough and flexible enough to allow the binding protein moiety and target peptide moiety to freely interact and respond to analyte concentration. In order to optimize the FRET effect, the average distance between the donor and acceptor fluorescent protein moieties should become between about 1 nm and about 10 nm, preferably between about 1 nm and about 6 nm, and more preferably between about 1 nm and about 4 nm, when the analyte is bound (or released). If the linker is too short or too stiff, the donor and acceptor protein moieties cannot readily change position. If the linker moiety is too long, the target peptide moiety might not bind to the binding protein moiety effectively. The linker moiety is, preferably, a peptide moiety. The preferred linker moiety is a peptide between about one and 30 amino acid residues in length, preferably between about two and 15 amino acid residues. One preferred linker moiety is a -Gly-Gly- linker.

The linker moiety can include flexible spacer amino acid sequences, such as those known in single-chain antibody research. For example, the linker moiety can be GGGGS (GGGGS)_n, GKSSGSGSESKS, GSTSGSGKSSEGKG, GSTSGSGKSSEGSSTKG, GSTSGSGKSSEGKG, GSTSGSGKPGSGEGSTKG, or EGKSSGSGSESKEF. Linking moieties are described, for example, in Huston, J.S., *et al.*, *PNAS* 85:5879-5883 (1988),

WO 98/40477

PCT/US98/04978

- 23 -

Whitlow, M., *et al.*, *Protein Engineering* 6:989-995 (1993), and Newton, D.L., *et al.*, *Biochemistry* 35:545-553 (1996).

Another embodiment of the invention includes a nucleic acid molecule, comprising a polynucleotide that encodes a fluorescent protein energy transfer partner, comprising: a
5 first chemical entity binding region that binds to a chemical entity and binds to a second protein partner, and a fluorescent protein in frame with the first chemical entity binding region; wherein the fluorescent protein is a first energy transfer partner for a second energy transfer partner attached to the second protein partner. Typically the fluorescent protein is an *Aequorea*-related fluorescent protein.

10 The first chemical entity binding region can comprise a bound conformation of smaller volume when the chemical entity is bound to the first chemical entity binding region compared to the first chemical entity binding region when the chemical entity is not bound to the first chemical entity binding region. The bound conformation can permit increased energy transfer between the first energy transfer partner and the second energy
15 transfer partner.

The bound conformation may also increase the affinity between the first chemical entity binding region and the second protein partner. Often the first chemical entity binding region binds to a protein association region of the second protein partner. In many cases the chemical entity is an analyte, as described herein. The chemical entity could also
20 include other entities such as cellular proteins.

Preferably, the first chemical entity binding region is either calmodulin, a calmodulin-related protein moiety, cGMP-dependent protein kinase, a steroid hormone receptor, a ligand binding domain of a steroid hormone receptor, protein kinase C, inositol-1,4,5-triphosphate receptor, or recoverin. More preferably, the first chemical entity binding
25 region is calmodulin or a calmodulin-related protein.

WO 98/40477

PCT/US98/04978

- 24 -

In some aspects of this embodiment of the invention the fluorescent protein energy transfer partner further comprises a second energy transfer partner in frame with the first energy transfer partner. Such embodiments can be produced by molecular biology techniques as opposed to protein synthesis techniques. Proteins can be produced by either technique to produce fluorescent protein that act as energy transfer partners. For example, the first energy transfer partner is either P4-3, EBFP, or W1B, and the second energy transfer partner is either S65T, EGFP, or 10c.

In addition energy partners can be generated using labelling methods known in the art for tagging proteins with fluorescent labels (e.g., biochemical reagents and non-protein fluorescent tags).

The fluorescent protein energy transfer partner can also further comprise a second protein partner comprising the second energy transfer partner in frame with the first energy transfer partner. The second protein partner can also further comprises a protein association region.

The second protein partner is either a calmodulin-binding domain of skMLCKp, smMLCK, CaMKII, Caldesmon, Calspermin, phosphofructokinase calcineurin, phosphorylase kinase, Ca²⁺-ATPase 59 kDa PDE, 60 kDa PDE, nitric oxide synthase, type I adenylyl cyclase, *Bordetella pertussis* adenylyl cyclase, Neuromodulin, Spectrin, MARCKS, F52, B-Adducin, HSP90a, HIV-1 gp160, BBMHBI, Dilute MHC, Mastoparan, Melittin, Glucagon, Secretin, VIP, GIP, or Model Peptide CBP2. Preferably, the protein association region is M13. The invention can include an amino acid linker in frame with the first chemical entity binding region and fusing the protein association region with the first chemical entity binding region. The invention can also include a fluorescent protein energy transfer partner that further comprises a localization sequence.

WO 98/40477

PCT/US98/04978

- 25 -

Another aspect of the present invention is a nucleic acid molecule, comprising a polynucleotide that encodes 1) a first fluorescent protein energy transfer partner, comprising: a chemical entity binding region that binds to a chemical entity and binds to a second fluorescent protein energy transfer partner, and a first fluorescent protein in
5 frame with the first chemical entity binding region; and 2) the second fluorescent protein energy transfer partner, comprising: a protein association region that binds to the chemical entity binding region, and a second fluorescent protein in frame with the protein association region; wherein the first fluorescent protein and the second fluorescent protein are energy transfer partners and the first fluorescent protein energy transfer
10 partner is in frame with the second first fluorescent protein energy transfer partner. The first and second fluorescent proteins can be *Aequorea*-related fluorescent proteins.

A further aspect of the present invention is this nucleic acid molecule, wherein the chemical entity binding region comprises a bound conformation of smaller volume when the chemical entity is bound to the chemical entity binding region compared to the
15 chemical entity binding region when the chemical entity is not bound to the chemical entity binding region and the bound conformation permits increased energy transfer between the first fluorescent protein energy transfer partner and the second fluorescent protein energy transfer partner. Another aspect of the present invention is this nucleic acid, wherein the bound conformation increases the affinity between the chemical entity
20 binding region and the second fluorescent protein energy transfer partner. The chemical entity is an analyte.

Another aspect of the present invention is a system for monitoring protein-protein association, comprising: 1) a first fluorescent protein energy transfer partner, comprising: a chemical entity binding region that binds to a chemical entity and binds to a second
25 fluorescent protein energy transfer partner, and a first energy transfer partner; and 2) the second fluorescent protein energy transfer partner, comprising: a protein association region that binds to the chemical entity binding region, and a second energy transfer partner; wherein the first energy transfer partner and the energy transfer partner are energy transfer partners.

WO 98/40477

PCT/US98/04978

- 26 -

The fluorescent indicators can also include a localization sequence to direct the indicator to particular cellular sites by fusion to appropriate organellar targeting signals or localized host proteins. A polynucleotide encoding a localization sequence, or signal sequence, can be ligated or fused at the 5' terminus of a polynucleotide encoding the
5 fluorescence indicator such that the signal peptide is located at the amino terminal end of the resulting fusion polynucleotide/polypeptide. In the case of eukaryotes, the signal peptide is believed to function to transport the fusion polypeptide across the endoplasmic reticulum. The secretory protein is then transported through the Golgi apparatus, into secretory vesicles and into the extracellular space or, preferably, the external
10 environment. Signal peptides which can be utilized according to the invention include pre-pro peptides which contain a proteolytic enzyme recognition site. Other signal peptides with similar properties to pro-calcitonin described herein are known to those skilled in the art, or can be readily ascertained without undue experimentation.

The localization sequence can be a nuclear localization sequence, an endoplasmic
15 reticulum localization sequence, a peroxisome localization sequence, a mitochondrial localization sequence, or a localized protein. Localization sequences can be targeting sequences which are described, for example, in "Protein Targeting", chapter 35 of Stryer, L., *Biochemistry* (4th ed.). W.H. Freeman, 1995. The localization sequence can also be a localized protein. Some important localization sequences include those targeting the
20 nucleus (KKKRRK), mitochondrion (amino terminal MLRTSSLFTRRVQPSLFRNLRQLQST-), endoplasmic reticulum (KDEL at C-terminus, assuming a signal sequence present at N-terminus), peroxisome (SKF at C-terminus), prenylation or insertion into plasma membrane (CaaX, CC, CXC, or CCXX at C-terminus), cytoplasmic side of plasma membrane (fusion to SNAP-25), or the Golgi
25 apparatus (fusion to furin).

The fluorescent indicators can be produced as fusion proteins by recombinant DNA technology. Recombinant production of fluorescent proteins involves expressing nucleic acids having sequences that encode the proteins. Nucleic acids encoding fluorescent proteins can be obtained by methods known in the art. For example, a nucleic acid

WO 98/40477

PCT/US98/04978

- 27 -

encoding the protein can be isolated by polymerase chain reaction of cDNA from *A. victoria* using primers based on the DNA sequence of *A. victoria* green fluorescent protein. PCR methods are described in, for example, U.S. Pat. No. 4,683,195; Mullis, *et al. Cold Spring Harbor Symp. Quant. Biol.* 51:263 (1987), and Erlich, ed., *PCR Technology*, (Stockton Press, NY, 1989). Mutant versions of fluorescent proteins can be made by site-specific mutagenesis of other nucleic acids encoding fluorescent proteins, or by random mutagenesis caused by increasing the error rate of PCR of the original polynucleotide with 0.1 mM MnCl₂ and unbalanced nucleotide concentrations. See, *e.g.*, U.S. patent application 08/337,915, filed November 10, 1994 or International application
10 PCT/US95/14692, filed 11/10/95.

The construction of expression vectors and the expression of genes in transfected cells involves the use of molecular cloning techniques also well known in the art. Sambrook *et al.*, *Molecular Cloning – A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, (1989) and *Current Protocols in Molecular Biology*, F.M. Ausubel
15 *et al.*, eds., (Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., most recent Supplement).

Nucleic acids used to transfect cells with sequences coding for expression of the polypeptide of interest generally will be in the form of an expression vector including expression control sequences operatively linked to a nucleotide sequence coding for
20 expression of the polypeptide. As used, the term "nucleotide sequence coding for expression of" a polypeptide refers to a sequence that, upon transcription and translation of mRNA, produces the polypeptide. This can include sequences containing, *e.g.*, introns. As used herein, the term "expression control sequences" refers to nucleic acid sequences that regulate the expression of a nucleic acid sequence to which it is
25 operatively linked. Expression control sequences are operatively linked to a nucleic acid sequence when the expression control sequences control and regulate the transcription and, as appropriate, translation of the nucleic acid sequence. Thus, expression control sequences can include appropriate promoters, enhancers, transcription terminators, a start codon (*i.e.*, ATG) in front of a protein-encoding gene, splicing signals for introns,

WO 98/40477

PCT/US98/04978

- 28 -

maintenance of the correct reading frame of that gene to permit proper translation of the mRNA, and stop codons.

Methods which are well known to those skilled in the art can be used to construct expression vectors containing the fluorescent indicator coding sequence and appropriate transcriptional/translational control signals. These methods include *in vitro* recombinant DNA techniques, synthetic techniques and *in vivo* recombination/genetic recombination. (See, for example, the techniques described in Maniatis, *et al.*, *Molecular Cloning A Laboratory Manual*, Cold Spring Harbor Laboratory, N.Y., 1989).

Transformation of a host cell with recombinant DNA may be carried out by conventional techniques as are well known to those skilled in the art. Where the host is prokaryotic, such as *E. coli*, competent cells which are capable of DNA uptake can be prepared from cells harvested after exponential growth phase and subsequently treated by the CaCl_2 method by procedures well known in the art. Alternatively, MgCl_2 or RbCl can be used. Transformation can also be performed after forming a protoplast of the host cell or by electroporation.

When the host is a eukaryote, such methods of transfection of DNA as calcium phosphate co-precipitates, conventional mechanical procedures such as microinjection, electroporation, insertion of a plasmid encased in liposomes, or virus vectors may be used. Eukaryotic cells can also be cotransfected with DNA sequences encoding the fusion polypeptide of the invention, and a second foreign DNA molecule encoding a selectable phenotype, such as the herpes simplex thymidine kinase gene. Another method is to use a eukaryotic viral vector, such as simian virus 40 (SV40) or bovine papilloma virus, to transiently infect or transform eukaryotic cells and express the protein. (*Eukaryotic Viral Vectors*, Cold Spring Harbor Laboratory, Gluzman ed., 1982). Preferably, a eukaryotic host is utilized as the host cell as described herein.

WO 98/40477

PCT/US98/04978

- 29 -

Techniques for the isolation and purification of either microbially or eukaryotically expressed polypeptides of the invention may be by any conventional means such as, for example, preparative chromatographic separations and immunological separations such as those involving the use of monoclonal or polyclonal antibodies or antigen.

- 5 A variety of host-expression vector systems may be utilized to express fluorescent indicator coding sequence. These include but are not limited to microorganisms such as bacteria transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing a fluorescent indicator coding sequence; yeast transformed with recombinant yeast expression vectors containing the fluorescent indicator coding sequence; plant cell systems infected with recombinant virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (*e.g.*, Ti plasmid) containing a fluorescent indicator coding sequence; insect cell systems infected with recombinant virus expression vectors (*e.g.*, baculovirus) containing a fluorescent indicator coding sequence; or animal cell systems infected with recombinant virus expression vectors (*e.g.*, retroviruses, adenovirus, vaccinia virus) containing a fluorescent indicator coding sequence, or transformed animal cell systems engineered for stable expression.

Depending on the host/vector system utilized, any of a number of suitable transcription and translation elements, including constitutive and inducible promoters, transcription enhancer elements, transcription terminators, *etc.* may be used in the expression vector (see, *e.g.*, Bitter, *et al.*, Methods in Enzymology 153:516-544, 1987). For example, when cloning in bacterial systems, inducible promoters such as pL of bacteriophage S, plac, ptrp, ptac (ptrp-lac hybrid promoter) and the like may be used. When cloning in mammalian cell systems, promoters derived from the genome of mammalian cells (*e.g.*, metallothionein promoter) or from mammalian viruses (*e.g.*, the retrovirus long terminal repeat; the adenovirus late promoter; the vaccinia virus 7.5K promoter) may be used. Promoters produced by recombinant DNA or synthetic techniques may also be used to provide for transcription of the inserted fluorescent indicator coding sequence.

WO 98/40477

PCT/US98/04978

- 30 -

In bacterial systems a number of expression vectors may be advantageously selected depending upon the use intended for the fluorescent indicator expressed. For example, when large quantities of the fluorescent indicator are to be produced, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Those which are engineered to contain a cleavage site to aid in recovering fluorescent indicator are preferred.

In yeast, a number of vectors containing constitutive or inducible promoters may be used. For a review see, Current Protocols in Molecular Biology, Vol. 2, Ed. Ausubel, *et al.*, Greene Publish. Assoc. & Wiley Interscience, Ch. 13, 1988; Grant, *et al.*, Expression and Secretion Vectors for Yeast, in Methods in Enzymology, Eds. Wu & Grossman, 31987, Acad. Press, N.Y., Vol. 153, pp.516-544, 1987; Glover, DNA Cloning, Vol. II, IRL Press, Wash., D.C., Ch. 3, 1986; and Bitter, Heterologous Gene Expression in Yeast, Methods in Enzymology, Eds. Berger & Kimmel, Acad. Press, N.Y., Vol. 152, pp. 673-684, 1987; and The Molecular Biology of the Yeast *Saccharomyces*, Eds. Strathern *et al.*, Cold Spring Harbor Press, Vols. I and II, 1982. A constitutive yeast promoter such as ADH or LEU2 or an inducible promoter such as GAL may be used (Cloning in Yeast, Ch. 3, R. Rothstein In: DNA Cloning Vol.11, A Practical Approach, Ed. DM Glover, IRL Press, Wash., D.C., 1986). Alternatively, vectors may be used which promote integration of foreign DNA sequences into the yeast chromosome.

In cases where plant expression vectors are used, the expression of a fluorescent indicator coding sequence may be driven by any of a number of promoters. For example, viral promoters such as the 35S RNA and 19S RNA promoters of CaMV (Brisson, *et al.*, *Nature* 310:511-514, 1984), or the coat protein promoter to TMV (Takamatsu, *et al.*, *EMBO J.* 6:307-311, 1987) may be used; alternatively, plant promoters such as the small subunit of RUBISCO (Coruzzi, *et al.*, 1984, *EMBO J.* 3:1671-1680; Broglie, *et al.*, *Science* 224:838-843, 1984); or heat shock promoters, *e.g.*, soybean hsp17.5-E or hsp17.3-B (Gurley, *et al.*, *Mol. Cell. Biol.* 6:559-565, 1986) may be used. These constructs can be introduced into plant cells using Ti plasmids, Ri plasmids, plant virus vectors,

WO 98/40477

PCT/US98/04978

- 31 -

direct DNA transformation, microinjection, electroporation, *etc.* For reviews of such techniques see, for example, Weissbach & Weissbach, *Methods for Plant Molecular Biology*, Academic Press, NY, Section VIII, pp. 421-463, 1988; and Grierson & Corey, *Plant Molecular Biology*, 2d Ed., Blackie, London, Ch. 7-9, 1988.

- 5 An alternative expression system which could be used to express fluorescent indicator is an insect system. In one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The fluorescent indicator coding sequence may be cloned
10 under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of the fluorescent indicator coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (*i.e.*, virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted gene is expressed,
15 see Smith, *et al.*, *J. Viol.* 46:584, 1983; Smith, U.S. Patent No. 4,215,051.

Eukaryotic systems, and preferably mammalian expression systems, allow for proper post-translational modifications of expressed mammalian proteins to occur. Eukaryotic cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, phosphorylation, and, advantageously secretion of the gene
20 product should be used as host cells for the expression of fluorescent indicator. Such host cell lines may include but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, Jurkat, HEK-293, and WI38.

Mammalian cell systems which utilize recombinant viruses or viral elements to direct expression may be engineered. For example, when using adenovirus expression vectors,
25 the fluorescent indicator coding sequence may be ligated to an adenovirus transcription/translation control complex, *e.g.*, the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (*e.g.*, region E1 or E3) will result in a recombinant virus that is viable and capable of

WO 98/40477

PCT/US98/04978

- 32 -

expressing the fluorescent indicator in infected hosts (e.g., see Logan & Shenk, *Proc. Natl. Acad. Sci. USA*, 81: 3655-3659, 1984). Alternatively, the vaccinia virus 7.5K promoter may be used. (e.g., see, Mackett, *et al.*, *Proc. Natl. Acad. Sci. USA*, 79: 7415-7419, 1982; Mackett, *et al.*, *J. Virol.* 49: 857-864, 1984; Panicali, *et al.*, *Proc. Natl. Acad. Sci. USA* 79: 4927-4931, 1982). Of particular interest are vectors based on bovine papilloma virus which have the ability to replicate as extrachromosomal elements (Sarver, *et al.*, *Mol. Cell. Biol.* 1: 486, 1981). Shortly after entry of this DNA into mouse cells, the plasmid replicates to about 100 to 200 copies per cell. Transcription of the inserted cDNA does not require integration of the plasmid into the host's chromosome, thereby yielding a high level of expression. These vectors can be used for stable expression by including a selectable marker in the plasmid, such as the *neo* gene. Alternatively, the retroviral genome can be modified for use as a vector capable of introducing and directing the expression of the fluorescent indicator gene in host cells (Cone & Mulligan, *Proc. Natl. Acad. Sci. USA*, 81:6349-6353, 1984). High level expression may also be achieved using inducible promoters, including, but not limited to, the metallothionine IIA promoter and heat shock promoters.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with the fluorescent indicator cDNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, *etc.*), and a selectable marker. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. For example, following the introduction of foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, *et al.*, *Cell*, 11: 223, 1977), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, *Proc. Natl. Acad. Sci. USA*, 48:2026, 1962), and adenine phosphoribosyltransferase (Lowy, *et al.*, *Cell*, 22: 817, 1980) genes

WO 98/40477

PCT/US98/04978

- 33 -

can be employed in tk⁻, hgp^rt⁻ or ap^rt⁻ cells respectively. Also, antimetabolite resistance can be used as the basis of selection for dhfr, which confers resistance to methotrexate (Wigler, *et al.*, *Proc. Natl. Acad. Sci. USA*, 77: 3567, 1980; O'Hare, *et al.*, *Proc. Natl. Acad. Sci. USA*, 8: 1527, 1981); gpt, which confers resistance to mycophenolic acid
5 (Mulligan & Berg, *Proc. Natl. Acad. Sci. USA*, 78: 2072, 1981; neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, *et al.*, *J. Mol. Biol.*, 150:1, 1981); and hyg^r, which confers resistance to hygromycin (Santerre, *et al.*, *Gene*, 30: 147, 1984) genes. Recently, additional selectable genes have been described, namely trpB, which allows cells to utilize indole in place of tryptophan; hisD, which allows cells
10 to utilize histinol in place of histidine (Hartman & Mulligan, *Proc. Natl. Acad. Sci. USA*, 85:8047, 1988); and ODC (ornithine decarboxylase) which confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine, DFMO (McConlogue L., In: *Current Communications in Molecular Biology*, Cold Spring Harbor Laboratory, ed., 1987).

15 DNA sequences encoding the fluorescence indicator polypeptide of the invention can be expressed *in vitro* by DNA transfer into a suitable host cell. "Host cells" are cells in which a vector can be propagated and its DNA expressed. The term also includes any progeny of the subject host cell. It is understood that all progeny may not be identical to the parental cell since there may be mutations that occur during replication. However,
20 such progeny are included when the term "host cell" is used. Methods of stable transfer, in other words when the foreign DNA is continuously maintained in the host, are known in the art.

Recombinant fluorescent protein can be produced by expression of nucleic acid encoding the protein in prokaryotes, such as *E. coli* or in eukaryotes, such as yeast cells or
25 mammalian cells. The fluorophore of *Aequorea*-related fluorescent proteins results from cyclization and oxidation of residues 65-67.

WO 98/40477

PCT/US98/04978

- 34 -

The construct can also contain a tag to simplify isolation of the fluorescent indicator. For example, a polyhistidine tag of, *e.g.*, six histidine residues, can be incorporated at the amino terminal end of the fluorescent protein. The polyhistidine tag allows convenient isolation of the protein in a single step by nickel-chelate chromatography.

- 5 In a preferred embodiment, the fluorescent indicator is a fusion protein produced by recombinant DNA technology in which a single polypeptide includes a donor moiety, a peptide linker moiety and an acceptor moiety. The donor moiety can be positioned at the amino-terminus relative to the acceptor moiety in the polypeptide. Such a fusion protein has the generalized structure: (amino terminus) donor fluorescent protein moiety
- 10 --peptide linker moiety --acceptor fluorescent protein moiety (carboxy terminus). Alternatively, the donor moiety can be positioned at the carboxy-terminus relative to the acceptor moiety within the fusion protein. Such a fusion protein has the generalized structure: (amino terminus) acceptor fluorescent protein moiety --peptide linker moiety --donor fluorescent protein moiety (carboxy terminus). The invention also envisions
- 15 fusion proteins that contain extra amino acid sequences at the amino and/or carboxy termini, for example, polyhistidine tags.

Thus, fluorescent indicators encoded by a recombinant nucleic acid include sequences coding for expression of a donor fluorescent protein moiety, an acceptor fluorescent protein moiety and a peptide linker moiety. The elements are selected so that upon

20 expression into a fusion protein, the donor and acceptor moieties exhibit FRET when the donor moiety is excited. The recombinant nucleic acid can be incorporated into an expression vector comprising expression control sequences operatively linked to the recombinant nucleic acid. The expression vector can be adapted for function in prokaryotes or eukaryotes by inclusion of appropriate promoters, replication sequences,

25 markers, *etc.*

The expression vector can be transfected into a host cell for expression of the recombinant nucleic acid. Host cells can be selected for high levels of expression in order to purify the fluorescent indicator fusion protein. *E. coli* is useful for this purpose.

WO 98/40477

PCT/US98/04978

- 35 -

Alternatively, the host cell can be a prokaryotic or eukaryotic cell selected to study the activity of an enzyme produced by the cell. In this case, the linker peptide is selected to include an amino acid sequence recognized by the protease. The cell can be, *e.g.*, a cultured cell or a cell *in vivo*.

- 5 A primary advantage of fluorescent indicator fusion proteins is that they are prepared by normal protein biosynthesis, thus completely avoiding organic synthesis and the requirement for customized unnatural amino acid analogs. The constructs can be expressed in *E. coli* in large scale for *in vitro* assays. Purification from bacteria is simplified when the sequences include polyhistidine tags for one-step purification by
10 nickel-chelate chromatography. Alternatively, the substrates can be expressed directly in a desired host cell for assays *in situ*.

In another embodiment, the invention provides a transgenic non-human animal that expresses a nucleic acid sequence which encodes the fluorescent indicator.

- The "non-human animals" of the invention comprise any non-human animal having
15 nucleic acid sequence which encodes a fluorescent indicator. Such non-human animals include vertebrates such as rodents, non-human primates, sheep, dog, cow, pig, amphibians, and reptiles. Preferred non-human animals are selected from the rodent family including rat and mouse, most preferably mouse. The "transgenic non-human animals" of the invention are produced by introducing "transgenes" into the germline of
20 the non-human animal. Embryonal target cells at various developmental stages can be used to introduce transgenes. Different methods are used depending on the stage of development of the embryonal target cell. The zygote is the best target for micro-injection. In the mouse, the male pronucleus reaches the size of approximately 20 micrometers in diameter which allows reproducible injection of 1-2 pl of DNA solution.
25 The use of zygotes as a target for gene transfer has a major advantage in that in most cases the injected DNA will be incorporated into the host gene before the first cleavage (Brinster *et al.*, *Proc. Natl. Acad. Sci. USA* 82:4438-4442, 1985). As a consequence, all cells of the transgenic non-human animal will carry the incorporated transgene. This will

WO 98/40477

PCT/US98/04978

- 36 -

in general also be reflected in the efficient transmission of the transgene to offspring of the founder since 50% of the germ cells will harbor the transgene. Microinjection of zygotes is the preferred method for incorporating transgenes in practicing the invention.

The term "transgenic" is used to describe an animal which includes exogenous genetic material within all of its cells. A "transgenic" animal can be produced by cross-breeding two chimeric animals which include exogenous genetic material within cells used in reproduction. Twenty-five percent of the resulting offspring will be transgenic *i.e.*, animals which include the exogenous genetic material within all of their cells in both alleles. 50% of the resulting animals will include the exogenous genetic material within one allele and 25% will include no exogenous genetic material.

Retroviral infection can also be used to introduce transgene into a non-human animal. The developing non-human embryo can be cultured *in vitro* to the blastocyst stage. During this time, the blastomeres can be targets for retro viral infection (Jaenich, R., *Proc. Natl. Acad. Sci USA* 73:1260-1264, 1976). Efficient infection of the blastomeres is obtained by enzymatic treatment to remove the zona pellucida (Hogan, *et al.* (1986) in *Manipulating the Mouse Embryo*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.). The viral vector system used to introduce the transgene is typically a replication-defective retro virus carrying the transgene (Jahner, *et al.*, *Proc. Natl. Acad. Sci. USA* 82:6927-6931, 1985; Van der Putten, *et al.*, *Proc. Natl. Acad. Sci USA* 82:6148-6152, 1985). Transfection is easily and efficiently obtained by culturing the blastomeres on a monolayer of virus-producing cells (Van der Putten, *supra*; Stewart, *et al.*, *EMBO J.* 6:383-388, 1987). Alternatively, infection can be performed at a later stage. Virus or virus-producing cells can be injected into the blastocoele (D. Jahner *et al.*, *Nature* 298:623-628, 1982). Most of the founders will be mosaic for the transgene since incorporation occurs only in a subset of the cells which formed the transgenic nonhuman animal. Further, the founder may contain various retro viral insertions of the transgene at different positions in the genome which generally will segregate in the offspring. In

WO 98/40477

PCT/US98/04978

- 37 -

addition, it is also possible to introduce transgenes into the germ line, albeit with low efficiency, by intrauterine retro viral infection of the midgestation embryo (D. Jahner *et al.*, *supra*).

A third type of target cell for transgene introduction is the embryonal stem cell (ES). ES
5 cells are obtained from pre-implantation embryos cultured *in vitro* and fused with
embryos (M. J. Evans *et al.* *Nature* 292:154-156, 1981; M.O. Bradley *et al.*, *Nature* 309:
255-258, 1984; Gossler, *et al.*, *Proc. Natl. Acad. Sci USA* 83: 9065-9069, 1986; and
Robertson *et al.*, *Nature* 322:445-448, 1986). Transgenes can be efficiently introduced
into the ES cells by DNA transfection or by retro virus-mediated transduction. Such
10 transformed ES cells can thereafter be combined with blastocysts from a nonhuman
animal. The ES cells thereafter colonize the embryo and contribute to the germ line of
the resulting chimeric animal. (For review see Jaenisch, R., *Science* 240: 1468-1474,
1988).

"Transformed" means a cell into which (or into an ancestor of which) has been
15 introduced, by means of recombinant nucleic acid techniques, a heterologous nucleic acid
molecule. "Heterologous" refers to a nucleic acid sequence that either originates from
another species or is modified from either its original form or the form primarily
expressed in the cell.

"Transgene" means any piece of DNA which is inserted by artifice into a cell, and
20 becomes part of the genome of the organism (*i.e.*, either stably integrated or as a stable
extrachromosomal element) which develops from that cell. Such a transgene may
include a gene which is partly or entirely heterologous (*i.e.*, foreign) to the transgenic
organism, or may represent a gene homologous to an endogenous gene of the organism.
Included within this definition is a transgene created by the providing of an RNA
25 sequence which is transcribed into DNA and then incorporated into the genome. The
transgenes of the invention include DNA sequences which encode which encodes the
fluorescent indicator which may be expressed in a transgenic non-human animal. The
term "transgenic" as used herein additionally includes any organism whose genome has

WO 98/40477

PCT/US98/04978

- 38 -

been altered by *in vitro* manipulation of the early embryo or fertilized egg or by any transgenic technology to induce a specific gene knockout. The term "gene knockout" as used herein, refers to the targeted disruption of a gene *in vivo* with complete loss of function that has been achieved by any transgenic technology familiar to those in the art.

5 In one embodiment, transgenic animals having gene knockouts are those in which the target gene has been rendered nonfunctional by an insertion targeted to the gene to be rendered non-functional by homologous recombination. As used herein, the term "transgenic" includes any transgenic technology familiar to those in the art which can produce an organism carrying an introduced transgene or one in which an endogenous

10 gene has been rendered non-functional or "knocked out."

Without further elaboration, it is believed that one skilled in the art can, based on the description herein, utilize the present invention to its fullest extent. All patents and publications cited herein are hereby incorporated by reference. A fluorescent indicator for Ca^{2+} was produced by sandwiching CaM-M13 fusion, described in Porumb, T., *et al.*,

15 *Prot. Engineering* 7:109-115 (1994), between a blue (P4-3) and a green (S65T) GFP mutant, as illustrated in FIG. 2a. The chimeric cDNA was cloned at *Bam*HI/*Eco*RI sites of pRSETB (Invitrogen), HIS is the amino-terminal tag peptide containing the polyhistidine sequence and XCaM is Xenopus calmodulin. Chimeric proteins incorporating a polyhistidine tag were expressed in *Escherichia coli*, purified by

20 nickel-chelate and size-exclusion chromatography, and their fluorescence characterized. Referring to FIG. 2a, the fluorescent CaM-based indicator ("cameleon-1") readily changes emission color by retracting and extending a long tongue (M13) into and out of the mouth of the CaM.

The amino acid composition of the boundary regions between the CaM-M13 hybrid and

25 GFPs can be important to optimize protein folding and the Ca^{2+} -sensitivity of FRET. One particularly sensitive indicator is shown in FIG. 2b. Referring to FIG. 2b, the amino acid and nucleotide sequences of the boundaries between P4-3 and CaM and between M13 and S65T in cameleon-1 are shown. Cameleon-1 has an 11 amino acid deletion at the C-terminus of P4-3 and a 5 amino acid deletion at the C-terminus of M13. Two

WO 98/40477

PCT/US98/04978

- 39 -

restriction sites *Sph*I and *Sac*I are underlined, which were utilized to connect the genes of P4-3 and S65T to the CaM-M13 gene, respectively. To facilitate the folding of the GFP that is fused to any other protein, a few glycine residues are usually inserted into the boundary. See, for example, Porumb, T., *et al.*, *Prot. Engineering* 7:109-115 (1994).

5 Further, glycine residues were not introduced so that P4-3 and S65T were fused rigidly to the CaM and M13, respectively. The rigid fusion leads to more effective transduction of the conformational change of CaM-M13, causing a greater change in FRET efficiency.

The fluorescent indicator was efficiently expressed and folded in bacteria and increased its ratio of UV-excited 510 nm to 445 nm emissions by 70% upon binding Ca^{2+} , as shown

10 in FIG. 3. The emission spectra of cameleon-1 were measured in 100 mM KCl, 20 mM MOPS, 1 mM EDTA, KOH to pH 7.20, before and after addition of 2 mM CaCl_2 to give 1 mM free Ca^{2+} . The Ca^{2+} binding to EDTA caused a local acidification of the solution, and a small fraction of the protein was denatured. Thus the spectrum after the Ca^{2+} addition dropped down slightly (compare with FIG. 4a). The decrease in blue and

15 increase in green emission indicated that Ca^{2+} increased the efficiency of FRET from P4-3 to S65T, consistent with the expected decrease in distance between the two ends of the protein. The Ca^{2+} response was fully reversible upon chelation of Ca^{2+} .

The Ca^{2+} -specificity of the response of cameleon-1 was examined. Mg^{2+} , pH, and ionic strength did not alter the emission spectra of either the Ca^{2+} -saturated and

20 Ca^{2+} -unsaturated forms. The emission spectra of saturated and unsaturated cameleon-1 were also not affected by hydrophobic proteins such as bovine serum albumin. Isolated CaM saturated with Ca^{2+} typically becomes sticky with hydrophobic amino acids exposed to the surface. The CaM in cameleon-1, on the other hand, appears to interact preferentially with its intramolecularly-adjacent M13 peptide. The self-contained nature

25 of the system minimizes the possibility that the protein might interact with endogenous CaM-binding sequences in eukaryotic cells.

WO 98/40477

PCT/US98/04978

- 40 -

The Ca^{2+} binding behavior of cameleon-1 was examined. The CaM-M13 hybrid protein without GFPs displayed a biphasic Ca^{2+} binding with two dissociation constants (80 nM and 2 μM). See, Porumb, T., *et al.*, *Prot. Engineering* 7:109-115 (1994). Titration experiments revealed that the emission ratio of cameleon-1 has a biphasic Ca^{2+} dependency, as shown in FIGS. 4a and 4b. FIG. 4a shows the change in emission spectrum of cameleon-1 on titration with Ca^{2+} when excited at 380 nm. The titration was done using KHHEDTA/KCaHEDTA solutions at pH 7.47. For clarity only two intermediate concentrations of Ca^{2+} are shown.

FIG. 4b shows Ca^{2+} titration curves of cameleon-1 (open circles) and cameleon-1/E104Q (solid circles). Data points were from 4 independent experiments at different pH using Ca^{2+} /EGTA and Ca^{2+} /HEEDTA systems for each protein. In each experiment, the emission ratio (510/445 nm) change was normalized to the value of full saturation with Ca^{2+} , which increased by 60-75% over the value of $^{2+}\text{zero Ca}$. The data of cameleon-1/E104Q were fitted to a four parameter logistic function curve (dotted line). The data of cameleon-1 were analysed using a linear combination of 2 four parameter logistic function fits (solid line). The apparent dissociation constants (K_d 's) for cameleon-1 were 68 nM and 11 μM , and the Hill coefficients were 1.8 and 1.0, respectively. The binding curve can be used to quantify the concentration of Ca^{2+} present in the sample. Because of simulated negative cooperativity, cameleon-1 covers a very wide range of Ca^{2+} concentration, from $<10^{-7}$ to $\sim 10^{-3}$ M.

The affinity of the CaM binding protein moiety can be modified. Many site-directed mutations have been studied for their effects on the Ca^{2+} binding and Ca^{2+} -induced conformational changes of CaM. See, Maune, J.F., *et al.*, *J. Biol. Chem.* 267:5286-5295 (1992), and Gao, Z.H., *et al.*, *J. Biol. Chem.* 268:20096-20104 (1993). For example, a mutant chimera protein with the conserved bidentate glutamic acid at position 12 of the third Ca^{2+} binding loop of the CaM mutated to glutamine (cameleon-1/E104Q) was constructed. The mutation eliminated the high-affinity response of cameleon-1, as indicated in FIG. 4b, (solid circles). Cameleon-1/E104Q showed a monophasic response (K_d , 4.4 μM ; Hill coefficient, 0.76), which corresponds closely to the low affinity

WO 98/40477

PCT/US98/04978

- 41 -

component of the cameleon-1 FRET response. Other modifications of CaM can be made to tune the Ca^{2+} affinities for particular applications of the fluorescent indicator.

HeLa cells were transfected with the recombinant plasmid (cameleon-1/pCDNA3) to determine whether cameleon-1 can work as a Ca^{2+} indicator in live cells. When the cells were excited with UV, however, the fluorescence of cameleon-1 was hardly detectable, mainly because of the dim fluorescence of the P4-3 component. Expression and folding of GFP at 37EC in mammalian cells was improved by introducing mammalian codon bias into the cDNA and mutating Phe64 to Leu, as in the commercially available construct "EGFP" (Clontech), which encodes F64L/S65T with mammalian codons.

10 The same changes were introduced into P4-3 (Y66H/Y145F), which did not change its final fluorescence properties but did improve expression in the HeLa cells. The improved blue mutant ("EBFP") and EGFP substituted P4-3 and S65T, respectively, in cameleon-1 to make cameleon-2, shown in FIG. 5, where Kz is Kozak's consensus sequence (M. Kozak, J. Cell Biol. 108:229-241 (1989). The nucleotide sequence (SEQ ID NO:1) and amino acid sequence (SEQ ID NO:2) of cameleon-2 are shown in FIG. 7.

15

The E104Q mutation which afforded low Ca^{2+} affinity in cameleon-1 was also introduced into cameleon-2; the resulting chimera protein (cameleon-2/E104Q) cameleon-3, also shown in FIG. 5. The nucleotide sequence (SEQ ID NO:5) and amino acid sequence (SEQ ID NO:6) of cameleon-3 are shown in FIG. 9.

20 Significant emission signals were observed from both the EBFP and EGFP in the cells expressing cameleon-2 or cameleon-3. Referring to FIG. 6a, fluorescence of cameleon-2 localized to the cytosol of HeLa cells. The fluorescence was uniformly distributed in the cytosolic compartment but excluded from the nucleus, as expected for a protein of 74 kDa without localization sequences or targeting signals. The image was taken using a 330WB80 (excitation filter) and a 535DF45 (emission filter). The bar is 10 μm .

25

FIG. 6b shows time courses of the spatially averaged green:blue emission ratios from two individual HeLa cells expressing cameleon-2. The two cells shown in FIG. 6a were

WO 98/40477

PCT/US98/04978

- 42 -

excited by UV (330WB80) and monitored every 15 seconds by digital imaging microscopy. See, for example, Tsien, R.Y. & Harootunian, A.T. *Cell Calcium*, 11:93-109 (1990). The emission bands (440DF40 and 535DF45) over the cytoplasmic regions were alternately sampled. *In situ* calibration was performed for each of the cells. The pre-stimulation ratio (arrowhead) was assumed to be 50 nM, and R_{\max} (arrow) the value after saturation with Ca^{2+} . The calculated values for pCas 7, 6, 5, 4 are indicated by horizontal bars on the right side of the panel.

Elevation of cytosolic Ca^{2+} concentration by saturating doses of ATP (as a purinergic agonist) and histamine produced significant increases in the emission ratio. Blockage of the histamine receptor by the antagonist cyproheptadine caused a rapid decrease in ratio, indicating the reversible behavior of the indicator. Addition of ionomycin followed by a high concentration (15 mM) of extracellular Ca^{2+} gave a large increase of the ratio (70-80% increase of the initial ratio value), which should correspond to the maximal ratio R_{\max} . Assuming the lowest ratio observed before stimulation represents the R_{\min} , calibration for free Ca^{2+} concentration can be performed. See, Adams, S.R., *et al.*, in *Fluorescent and Luminescent Probes for Biological Activity* (ed. Mason, W.T.) (Academic Press, 1993).

By contrast, cameleon-3, which lacks the high affinity component of Ca^{2+} binding to CaM, did not detect the changes in cytosolic Ca^{2+} concentration signals due to ATP or histamine, but gave a similar R_{\max} in response to ionomycin and 20 mM extracellular Ca^{2+} . Cameleon-3 is less sensitive to and buffers cytosolic Ca^{2+} to a lesser extent than does cameleon-2. The probable high Ca^{2+} dissociation rate of cameleon-3 is advantageous for tracking rapid Ca^{2+} release kinetics. The *in vitro* study revealed that the cameleon indicators show a relatively fast cellular response to Ca^{2+} concentration changes.

Addition of a nuclear localization sequence to cameleon-2 yielded a Ca^{2+} indicator, cameleon-2nu, shown in FIG. 5. The nucleotide sequence (SEQ ID NO:3) and amino acid sequence (SEQ ID NO:4) of cameleon-2nu are shown in FIG. 8. The fluorescence

WO 98/40477

PCT/US98/04978

- 43 -

of cameleon-2nu was localized to nuclei, as depicted in FIG. 6c. The time course of Ca^{2+} concentrations in nuclei was followed (FIG. 6d) and was similar to the results obtained in the cytosol (FIG. 6b). Agonist-induced changes in the free Ca^{2+} concentration inside the endoplasmic reticulum in intact cells were similarly monitored. The low-affinity indicator cameleon-3 was engineered to reside in the lumen of endoplasmic reticulum (ER) (cameleon-3er) (FIG. 5) by addition of a localization sequence at the amino terminus and a KDEL signal for ER retention at the carboxy terminus of the fluorescent indicator. The nucleotide sequence (SEQ ID NO:7) and amino acid sequence of cameleon-3er (SEQ ID NO:8) are shown in FIG. 10. Reticular patterns of fluorescence were seen in HeLa cells expressing the protein. FIG. 6e is a digital fluorescence image of cameleon-3er in transfected HeLa cells. The image was obtained with a cooled CCD camera system with a 480DF30 (excitation filter) and a 535DF45 (emission filter). The bar is 10 μm .

FIG. 6f shows a time course of the average $\text{Ca}^{2+}_{\text{er}}$ concentration of four cells obtained with a video-rate confocal microscope. Digital fluorescence images were the result of simultaneous acquisition of two confocal single-wavelength emission images at 450 nm (65 nm bandpass) and 535 nm (45 nm bandpass). After background subtraction, the ratio of the long wavelength image over the short wavelength one was calculated. Cells were illuminated for 66 msec (2 frames) for each time point. The interference filters and dichroics, and the sensitivity of the detectors are different between the CCD and confocal microscope systems. Therefore, the ratios obtained in the two systems differ quantitatively.

The pre-stimulus $\text{Ca}^{2+}_{\text{er}}$ concentration reported by cameleon-3er was consistently higher than cytosolic or nuclear Ca^{2+} concentrations reported by cameleon-2 and cameleon-2nu, respectively. Histamine reproducibly decreased the $\text{Ca}^{2+}_{\text{er}}$ concentration in all of 15 cells, whereas it always increased the cytosolic and nuclear Ca^{2+} concentrations. Receptor blockade by cyproheptadine reversed the decrease $\text{Ca}^{2+}_{\text{er}}$ concentration, indicating refill of the Ca^{2+} pools. In FIG. 6f, the ratio did not reach the value of the resting state, whereas complete reversion was observed in five other experiments. Addition of

WO 98/40477

PCT/US98/04978

- 44 -

ionomycin and 20 mM extracellular Ca^{2+} increased $\text{Ca}^{2+}_{\text{er}}$ concentration to a saturating value above the starting level. Changes in $\text{Ca}^{2+}_{\text{er}}$ concentration were generally slower than those of the cytosol and nuclei.

The targetability of the fluorescent indicators can permit Ca^{2+} measurements at previously inaccessible sites such as the immediate vicinity of synaptic vesicles or Ca^{2+} channels, and in genetically tractable organisms that do not load well with esters of small-molecule indicators.

The Examples described below are illustrative of the disclosed method; however, many alternatives, modifications and variations will be clear to those skilled in the art.

10 Examples

Gene construction

The cDNA of the GFP mutant P4-3 was amplified by PCR with a sense primer containing a *Bam*HI site, and a reverse primer containing an *Sph*I site and eliminating the GFP stop codon. See, for example, Heim, R. & Tsien, R.Y. *Current Biol.* 6:178-182 (1996). Likewise, the cDNA of S65T was amplified with a *Sac*I site and an *Eco*RI site introduced to the 5' and 3' ends of the gene, respectively. Two restriction sites (*Sph*I and *Sac*I) were introduced by PCR into 5' and 3' ends of the CaM-M13 gene, respectively, using the pHY1 as a template. See, Porumb, T., *et al.* *Prot. Engineering* 7:109-115 (1994). All the amplification reactions were done by Pfu polymerase (Stratagene). The restricted products were ligated and cloned in-frame into the *Bam*HI/*Eco*RI sites of pRSETB (Invitrogen). The modifications of the boundary regions between P4-3 and CaM and between M13 and S65T were performed by PCR or by a combined use of restriction enzymes, Klenow fragment of DNA polymerase I, T4 DNA polymerase, mung bean exonuclease, and T4 DNA ligase as described, for example, in *Molecular Cloning, A Laboratory Manual* (eds. Sambrook, J., Fritsch, E.F. & Maniatis, T.) (CSH Laboratory Press, 1989). The pHGFP plasmid was commercially available from Clontech. Two amino acid substitutions (Y66H and Y145F) were made in hEGFP to construct EBFP. Oligonucleotide-directed mutageneses were carried out using the Muta-Gene Phagemid

WO 98/40477

PCT/US98/04978

- 45 -

in vitro kit (Bio-Rad) at the codons for Y66H and Y145F of EGFP, and for E104Q of the calmodulin. The 5' end of the EBFP gene was modified by PCR to have a *HindIII* restriction site followed by a Kozak's consensus sequence (ACCGCC-ATG). The *HindIII/EcoRI* fragment encoding the entire chimeric protein was subcloned in the
5 mammalian expression vector pCDNA3 (Invitrogen).

For cameleon-2nu, the cameleon-2 DNA was extended by PCR at the 3' end with the sequence encoding the nuclear localization sequence (PKKKRKVEDP). See, Forbes, D.J. *Ann.Rev.Cell Biol.* 8:495-527 (1992). Cameleon-3er DNA was likewise obtained by extending the cameleon-3 DNA at the 5' end with the sequence encoding the
10 localization sequence peptide from calreticulin (MLLPVPLLLGLLGLAAAD), and at the 3' end with the sequence encoding the ER retention signal (KDEL). See, Kendall, J.M. *et al., Biochem. Biophys. Res. Commun.* 189:1008-1016 (1992).

Protein expression and spectroscopy

The expression of chimera proteins in bacteria was performed using the T7 expresson
15 system (pRSETB/JM109(DE3)). Cultures were grown at room temperature, and protein expression was induced by isopropyl B-D-thiogalactoside. Cells were lysed by a French press. The polyhistidine-tagged chimera proteins were purified from the cleared lysates on nickel-chelate columns (Qiagen). The protein samples in the eluates were concentrated by Centricon 30 (Amicon), and were further purified by gel-filtration
20 column to remove abortive chimera proteins which resulted from proteolysis or misfolding. Emission spectra of the purified proteins were measured using a fluorometer (Spex Industries, Edison, NJ) at excitation 380 nm.

Ca²⁺ titration and calibration

The titration experiments were performed by the "pH-metric method" as described in
25 Grzegorz, G., *et al., J. Biol. Chem.* 260:3440-3450 (1985). *In situ* calibration for cytosolic Ca²⁺ concentration utilized the equation:

$$[Ca^{2+}]_c = K'_d ((R - R_{min}) / (R_{max} - R))^{(1/nH)}$$

WO 98/40477

PCT/US98/04978

- 46 -

where K'_d is the apparent dissociation constant corresponding to the Ca^{2+} concentration at which R is midway between R_{\max} and R_{\min} , and nH is the Hill coefficient.

Imaging

Two to five days after the cDNA transfection with lipofectin (Gibco BRL), the cells were
5 imaged on a Zeiss Axiovert microscope with a cooled CCD camera (Photometrics, Tucson, AZ) interfaced to a personal computer. The program MetaFluor 2.75 (Universal Imaging) was used for controlling data acquisition and analysis. Dual-emission ratio imaging was carried out by manually switching the two emission filters (440DF40 for EBFP, 535DF45 for EGFP) in front of a single imaging camera. The excitation filter
10 (330WB80) was used with a 420DRLP dichroic mirror. Digital fluorescence imaging with a video-rate confocal microscope was performed as described in Tsien, R.Y. & Backskai, B.J. *Handbook of Biological Confocal Microscopy* (ed. Pawley, J.B.) (Plenum Press, New York, 1995) p. 459-478. Cells were illuminated with wide band UV (351-364 nm) from an Ar^+ ion laser. The primary dichroic (380DRLP) reflects UV and
15 transmits light emitted from the specimen, which is subsequently split by a secondary dichroic (505DRLP) into two broad bands: EBFP emission (450DF65) and EGFP emission (535DF45), and counted by photomultiplier tubes.

WO 98/40477

PCT/US98/04978

- 47 -

SEQUENCE LISTING

- (1) GENERAL INFORMATION:
- (i) APPLICANT: The Regents of the University of California
- (ii) TITLE OF INVENTION: Fluorescent Protein Sensors for Detection of Analytes
- (iii) NUMBER OF SEQUENCES: 8
- (iv) CORRESPONDENCE ADDRESS:
- (A) ADDRESSEE: Fish & Richardson P.C.
- (B) STREET: 4225 Executive Square, Suite 1400
- (C) CITY: La Jolla
- (D) STATE: CA
- (E) COUNTRY: USA
- (F) ZIP: 92037
- (v) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: US 08/818,252
- (B) FILING DATE: 14 March 1997
- (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER US 08/818,253
- (B) FILING DATE: 14 MARCH 1997
- (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: US 08/919,143
- (B) FILING DATE: 27 AUGUST 1997
- (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
- (A) NAME: Haile, Lisa A.
- (B) REGISTRATION NUMBER: 38,347
- (C) REFERENCE/DOCKET NUMBER: 07257/058WO1
- (ix) TELECOMMUNICATION INFORMATION:
- (A) TELEPHONE: 619/678-5070
- (B) TELEFAX: 619/678-5099
- (2) INFORMATION FOR SEQ ID NO:1:
- (i) SEQUENCE CHARACTERISTICS
- (A) LENGTH: 1929 BASE PAIRS
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: cDNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATG GTG AGC AAG GGC GAG GAG CTG TTC ACC GGG GTG GTG CCC ATC CTG 48
Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu
5 10 15

WO 98/40477

PCT/US98/04978

- 48 -

GTC	GAG	CTG	GAC	GGC	GAC	GTA	AAC	GGC	CAC	AAG	TTC	AGC	GTG	TCC	GGC	96
Val	Glu	Leu	Asp	Gly	Asp	Val	Asn	Gly	His	Lys	Phe	Ser	Val	Ser	Gly	
			20					25					30			
GAG	GGC	GAG	GGC	GAT	GCC	ACC	TAC	GGC	AAG	CTG	ACC	CTG	AAG	TTC	ATC	144
Glu	Gly	Glu	Gly	Asp	Ala	Thr	Tyr	Gly	Lys	Leu	Thr	Leu	Lys	Phe	Ile	
		35					40					45				
TGC	ACC	ACC	GGC	AAG	CTG	CCC	GTG	CCC	TGG	CCC	ACC	CTC	GTG	ACC	ACC	192
Cys	Thr	Thr	Gly	Lys	Leu	Pro	Val	Pro	Trp	Pro	Thr	Leu	Val	Thr	Thr	
	50					55					60					
CTG	ACC	CAT	GGC	GTG	CAG	TGC	TTC	AGC	CGC	TAC	CCC	GAC	CAC	ATG	AAG	240
Leu	Thr	His	Gly	Val	Gln	Cys	Phe	Ser	Arg	Tyr	Pro	Asp	His	Met	Lys	
	65				70				75						80	
CAG	CAC	GAC	TTC	TTC	AAG	TCC	GCC	ATG	CCC	GAA	GGC	TAC	GTG	CAG	GAG	288
Gln	His	Asp	Phe	Phe	Lys	Ser	Ala	Met	Pro	Glu	Gly	Tyr	Val	Gln	Glu	
				85				90						95		
CGC	ACC	ATC	TTC	TTC	AAG	GAC	GAC	GGC	AAC	TAC	AAG	ACC	CGC	GCC	GAG	336
Arg	Thr	Ile	Phe	Phe	Lys	Asp	Asp	Gly	Asn	Tyr	Lys	Thr	Arg	Ala	Glu	
			100					105					110			
GTG	AAG	TTC	GAG	GGC	GAC	ACC	CTG	GTG	AAC	CGC	ATC	GAG	CTG	AAG	GGC	384
Val	Lys	Phe	Glu	Gly	Asp	Thr	Leu	Val	Asn	Arg	Ile	Glu	Leu	Lys	Gly	
		115					120					125				
ATC	GAC	TTC	AAG	GAG	GAC	GGC	AAC	ATC	CTG	GGG	CAC	AAG	CTG	GAG	TAC	432
Ile	Asp	Phe	Lys	Glu	Asp	Gly	Asn	Ile	Leu	Gly	His	Lys	Leu	Glu	Tyr	
	130					135				140						
AAC	TTC	AAC	AGC	CAC	AAC	GTC	TAT	ATC	ATG	GCC	GAC	AAG	CAG	AAG	AAC	480
Asn	Phe	Asn	Ser	His	Asn	Val	Tyr	Ile	Met	Ala	Asp	Lys	Gln	Lys	Asn	
	145				150					155					160	
GGC	ATC	AAG	GTG	AAC	TTC	AAG	ATC	CGC	CAC	AAC	ATC	GAG	GAC	GGC	AGC	528
Gly	Ile	Lys	Val	Asn	Phe	Lys	Ile	Arg	His	Asn	Ile	Glu	Asp	Gly	Ser	
				165				170						175		
GTG	CAG	CTC	GCC	GAC	CAC	TAC	CAG	CAG	AAC	ACC	CCC	ATC	GGC	GAC	GGC	576
Val	Gln	Leu	Ala	Asp	His	Tyr	Gln	Gln	Asn	Thr	Pro	Ile	Gly	Asp	Gly	
			180					185					190			
CCC	GTG	CTG	CTG	CCC	GAC	AAC	CAC	TAC	CTG	AGC	ACC	CAG	TCC	GCC	CTG	624
Pro	Val	Leu	Leu	Pro	Asp	Asn	His	Tyr	Leu	Ser	Thr	Gln	Ser	Ala	Leu	
		195					200					205				
AGC	AAA	GAC	CCC	AAC	GAG	AAG	CGC	GAT	CAC	ATG	GTC	CTG	CTG	GAG	TTC	672
Ser	Lys	Asp	Pro	Asn	Glu	Lys	Arg	Asp	His	Met	Val	Leu	Leu	Glu	Phe	
	210					215					220					
GTG	ACC	GCC	GCC	CGC	ATG	CAT	GAC	CAA	CTG	ACA	GAA	GAG	CAG	ATT	GCA	720
Val	Thr	Ala	Ala	Arg	Met	His	Asp	Gln	Leu	Thr	Glu	Glu	Gln	Ile	Ala	
	225				230					235					240	
GAG	TTC	AAA	GAA	GCC	TTC	TCA	TTA	TTC	GAC	AAG	GAT	GGG	GAC	GGC	ACC	768
Glu	Phe	Lys	Glu	Ala	Phe	Ser	Leu	Phe	Asp	Lys	Asp	Gly	Asp	Gly	Thr	
				245				250						255		
ATC	ACC	ACA	AAG	GAA	CTT	GGC	ACC	GTT	ATG	AGG	TCG	CTT	GGA	CAA	AAC	816
Ile	Thr	Thr	Lys	Glu	Leu	Gly	Thr	Val	Met	Arg	Ser	Leu	Gly	Gln	Asn	
			260					265					270			

WO 98/40477

PCT/US98/04978

- 49 -

CCA	ACG	GAA	GCA	GAA	TTG	CAG	GAT	ATG	ATC	AAT	GAA	GTC	GAT	GCT	GAT	864
Pro	Thr	Glu	Ala	Glu	Leu	Gln	Asp	Met	Ile	Asn	Glu	Val	Asp	Ala	Asp	
		275					280					285				
GGC	AAT	GGA	ACG	ATT	TAC	TTT	CCT	GAA	TTT	CTT	ACT	ATG	ATG	GCT	AGA	912
Gly	Asn	Gly	Thr	Ile	Tyr	Phe	Pro	Glu	Phe	Leu	Thr	Met	Met	Ala	Arg	
	290					295					300					
AAA	ATG	AAG	GAC	ACA	GAC	AGC	GAA	GAG	GAA	ATC	CGA	GAA	GCA	TTC	CGT	960
Lys	Met	Lys	Asp	Thr	Asp	Ser	Glu	Glu	Glu	Ile	Arg	Glu	Ala	Phe	Arg	
305					310					315					320	
GTT	TTT	GAC	AAG	GAT	GGG	AAC	GGC	TAC	ATC	AGC	GCT	GCT	GAA	TTA	CGT	1008
Val	Phe	Asp	Lys	Asp	Gly	Asn	Gly	Tyr	Ile	Ser	Ala	Ala	Glu	Leu	Arg	
				325					330					335		
CAC	GTC	ATG	ACA	AAC	CTC	GGG	GAG	AAG	TTA	ACA	GAT	GAA	GAA	GTT	GAT	1056
His	Val	Met	Thr	Asn	Leu	Gly	Glu	Lys	Leu	Thr	Asp	Glu	Glu	Val	Asp	
			340					345					350			
GAA	ATG	ATA	AGG	GAA	GCA	GAT	ATC	GAT	GGT	GAT	GGC	CAA	GTA	AAC	TAT	1104
Glu	Met	Ile	Arg	Glu	Ala	Asp	Ile	Asp	Gly	Asp	Gly	Gln	Val	Asn	Tyr	
		355					360					365				
GAA	GAG	TTT	GTA	CAA	ATG	ATG	ACA	GCA	AAG	GGG	GGG	AAG	AGG	CGC	TGG	1152
Glu	Glu	Phe	Val	Gln	Met	Met	Thr	Ala	Lys	Gly	Gly	Lys	Arg	Arg	Trp	
	370					375					380					
AAG	AAA	AAC	TTC	ATT	GCC	GTC	AGC	GCT	GCC	AAC	CGG	TTC	AAG	AAG	ATC	1200
Lys	Lys	Asn	Phe	Ile	Ala	Val	Ser	Ala	Ala	Asn	Arg	Phe	Lys	Lys	Ile	
385					390					395					400	
TCC	GAG	CTC	ATG	GTG	AGC	AAG	GGC	GAG	GAG	CTG	TTC	ACC	GGG	GTG	GTG	1248
Ser	Glu	Leu	Met	Val	Ser	Lys	Gly	Glu	Glu	Leu	Phe	Thr	Gly	Val	Val	
				405					410					415		
CCC	ATC	CTG	GTC	GAG	CTG	GAC	GGC	GAC	GTA	AAC	GGC	CAC	AAG	TTC	AGC	1296
Pro	Ile	Leu	Val	Glu	Leu	Asp	Gly	Asp	Val	Asn	Gly	His	Lys	Phe	Ser	
			420					425					430			
GTG	TCC	GGC	GAG	GGC	GAG	GGC	GAT	GCC	ACC	TAC	GGC	AAG	CTG	ACC	CTG	1344
Val	Ser	Gly	Glu	Gly	Glu	Gly	Asp	Ala	Thr	Tyr	Gly	Lys	Leu	Thr	Leu	
		435					440					445				
AAG	TTC	ATC	TGC	ACC	ACC	GGC	AAG	CTG	CCC	GTG	CCC	TGG	CCC	ACC	CTC	1392
Lys	Phe	Ile	Cys	Thr	Thr	Gly	Lys	Leu	Pro	Val	Pro	Trp	Pro	Thr	Leu	
	450					455					460					
GTG	ACC	ACC	CTG	ACC	TAC	GGC	GTG	CAG	TGC	TTC	AGC	CGC	TAC	CCC	GAC	1440
Val	Thr	Thr	Leu	Thr	Tyr	Gly	Val	Gln	Cys	Phe	Ser	Arg	Tyr	Pro	Asp	
	465				470				475						480	
CAC	ATG	AAG	CAG	CAC	GAC	TTC	TTC	AAG	TCC	GCC	ATG	CCC	GAA	GGC	TAC	1488
His	Met	Lys	Gln	His	Asp	Phe	Phe	Lys	Ser	Ala	Met	Pro	Glu	Gly	Tyr	
			485					490						495		
GTC	CAG	GAG	CGC	ACC	ATC	TTC	TTC	AAG	GAC	GAC	GGC	AAC	TAC	AAG	ACC	1536
Val	Gln	Glu	Arg	Thr	Ile	Phe	Phe	Lys	Asp	Asp	Gly	Asn	Tyr	Lys	Thr	
			500					505					510			
CGC	GCC	GAG	GTG	AAG	TTC	GAG	GGC	GAC	ACC	CTG	GTG	AAC	CGC	ATC	GAG	1584
Arg	Ala	Glu	Val	Lys	Phe	Glu	Gly	Asp	Thr	Leu	Val	Asn	Arg	Ile	Glu	
		515					520						525			

WO 98/40477

PCT/US98/04978

- 50 -

CTG AAG GGC ATC GAC TTC AAG GAG GAC GGC AAC ATC CTG GGG CAC AAG	1632
Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys	
530 535 540	
CTG GAG TAC AAC TAC AAC AGC CAC AAC GTC TAT ATC ATG GCC GAC AAG	1680
Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys	
545 550 555 560	
CAG AAG AAC GGC ATC AAG GTG AAC TTC AAG ATC CGC CAC AAC ATC GAG	1728
Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu	
565 570 575	
GAC GGC AGC GTG CAG CTC GCC GAC CAC TAC CAG CAG AAC ACC CCC ATC	1776
Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile	
580 585 590	
GGC GAC GGC CCC GTG CTG CTG CCC GAC AAC CAC TAC CTG AGC ACC CAG	1824
Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln	
595 600 605	
TCC GCC CTG AGC AAA GAC CCC AAC GAG AAG CGC GAT CAC ATG GTC CTG	1872
Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu	
610 615 620	
CTG GAG TTC GTG ACC GCC GCC GGG ATC ACT CTC GGC ATG GAC GAG CTG	1920
Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu	
625 630 635 640	
TAC AAG TAA	1929
Tyr Lys	

(3) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS
 (A) LENGTH: 642 AMINO ACIDS
 (B) TYPE: AMINO ACID
 (C) STRANDEDNESS:
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu	5 10 15
Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly	20 25 30
Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile	35 40 45
Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr	50 55 60
Leu Thr His Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys	65 70 75 80
Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu	85 90 95
Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu	100 105 110
Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly	115 120 125

WO 98/40477

PCT/US98/04978

- 51 -

Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr
130 135 140
Asn Phe Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn
145 150 155 160
Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser
165 170 175
Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly
180 185 190
Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu
195 200 205
Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe
210 215 220
Val Thr Ala Ala Arg Met His Asp Gln Leu Thr Glu Glu Gln Ile Ala
225 230 235 240
Glu Phe Lys Glu Ala Phe Ser Leu Phe Asp Lys Asp Gly Asp Gly Thr
245 250 255
Ile Thr Thr Lys Glu Leu Gly Thr Val Met Arg Ser Leu Gly Gln Asn
260 265 270
Pro Thr Glu Ala Glu Leu Gln Asp Met Ile Asn Glu Val Asp Ala Asp
275 280 285
Gly Asn Gly Thr Ile Tyr Phe Pro Glu Phe Leu Thr Met Met Ala Arg
290 295 300
Lys Met Lys Asp Thr Asp Ser Glu Glu Glu Ile Arg Glu Ala Phe Arg
305 310 315 320
Val Phe Asp Lys Asp Gly Asn Gly Tyr Ile Ser Ala Ala Glu Leu Arg
325 330 335
His Val Met Thr Asn Leu Gly Glu Lys Leu Thr Asp Glu Glu Val Asp
340 345 350
Glu Met Ile Arg Glu Ala Asp Ile Asp Gly Asp Gly Gln Val Asn Tyr
355 360 365
Glu Glu Phe Val Gln Met Met Thr Ala Lys Gly Gly Lys Arg Arg Trp
370 375 380
Lys Lys Asn Phe Ile Ala Val Ser Ala Ala Asn Arg Phe Lys Lys Ile
385 390 395 400
Ser Glu Leu Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val
405 410 415
Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser
420 425 430
Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu
435 440 445
Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu
450 455 460
Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp
465 470 475 480

PCT/US98/04978

[illegible]

(i) SEQUENCE CHARACTERISTICS
(A) LENGTH: 1959 BASE PAIRS
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: SINGLE
(D) TOPOLOGY: LINEAR

(xi) SEQUENCE DESCRIPTION: SEO ID NO:3:

ATG	GTG	AGC	AAG	GGC	GAG	GAG	CTG	TTC	ACC	GGG	GTG	GTG	CCC	ATC	CTG	48
Met	Val	Ser	Lys	Gly	Glu	Glu	Leu	Phe	Thr	Gly	Val	Val	Pro	Ile	Leu	
				5					10					15		
GTC	GAG	CTG	GAC	GGC	GAC	GTA	AAC	GGC	CAC	AAG	TTC	AGC	GTG	TCC	GGC	96
Val	Glu	Leu	Asp	Gly	Asp	Val	Asn	Gly	His	Lys	Phe	Ser	Val	Ser	Gly	
			20					25					30			
GAG	GGC	GAG	GGC	GAT	GCC	ACC	TAC	GGC	AAG	CTG	ACC	CTG	AAG	TTC	ATC	
	144															
Glu	Gly	Glu	Gly	Asp	Ala	Thr	Tyr	Gly	Lys	Leu	Thr	Leu	Lys	Phe	Ile	
		35					40					45				
TGC	ACC	ACC	GGC	AAG	CTG	CCC	GTG	CCC	TGG	CCC	ACC	CTC	GTG	ACC	ACC	192
Cys	Thr	Thr	Gly	Lys	Leu	Pro	Val	Pro	Trp	Pro	Thr	Leu	Val	Thr	Thr	
	50					55					60					

WO 98/40477

PCT/US98/04978

- 53 -

CTG ACC CAT GGC GTG CAG TGC TTC AGC CGC TAC CCC GAC CAC ATG AAG	240
Leu Thr His Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys	
65 70 75 80	
CAG CAC GAC TTC TTC AAG TCC GCC ATG CCC GAA GGC TAC GTC CAG GAG	288
Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu	
85 90 95	
CGC ACC ATC TTC TTC AAG GAC GAC GGC AAC TAC AAG ACC CGC GCC GAG	336
Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu	
100 105 110	
GTG AAG TTC GAG GGC GAC ACC CTG GTG AAC CGC ATC GAG CTG AAG GGC	384
Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly	
115 120 125	
ATC GAC TTC AAG GAG GAC GGC AAC ATC CTG GGG CAC AAG CTG GAG TAC	432
Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr	
130 135 140	
AAC TTC AAC AGC CAC AAC GTC TAT ATC ATG GCC GAC AAG CAG AAG AAC	480
Asn Phe Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn	
145 150 155 160	
GGC ATC AAG GTG AAC TTC AAG ATC CGC CAC AAC ATC GAG GAC GGC AGC	528
Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser	
165 170 175	
GTG CAG CTC GCC GAC CAC TAC CAG CAG AAC ACC CCC ATC GGC GAC GGC	576
Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly	
180 185 190	
CCC GTG CTG CTG CCC GAC AAC CAC TAC CTG AGC ACC CAG TCC GCC CTG	624
Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu	
195 200 205	
AGC AAA GAC CCC AAC GAG AAG CGC GAT CAC ATG GTC CTG CTG GAG TTC	672
Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe	
210 215 220	
GTG ACC GCC GCC CGC ATG CAT GAC CAA CTG ACA GAA GAG CAG ATT GCA	720
Val Thr Ala Ala Arg Met His Asp Gln Leu Thr Glu Glu Gln Ile Ala	
225 230 235 240	
GAG TTC AAA GAA GCC TTC TCA TTA TTC GAC AAG GAT GGG GAC GGC ACC	768
Glu Phe Lys Glu Ala Phe Ser Leu Phe Asp Lys Asp Gly Asp Gly Thr	
245 250 255	
ATC ACC ACA AAG GAA CTT GGC ACC GTT ATG AGG TCG CTT GGA CAA AAC	816
Ile Thr Thr Lys Glu Leu Gly Thr Val Met Arg Ser Leu Gly Gln Asn	
260 265 270	
CCA ACG GAA GCA GAA TTG CAG GAT ATG ATC AAT GAA GTC GAT GCT GAT	864
Pro Thr Glu Ala Glu Leu Gln Asp Met Ile Asn Glu Val Asp Ala Asp	
275 280 285	
GGC AAT GGA ACG ATT TAC TTT CCT GAA TTT CTT ACT ATG ATG GCT AGA	912
Gly Asn Gly Thr Ile Tyr Phe Pro Glu Phe Leu Thr Met Met Ala Arg	
290 295 300	
AAA ATG AAG GAC ACA GAC AGC GAA GAG GAA ATC CGA GAA GCA TTC CGT	960
Lys Met Lys Asp Thr Asp Ser Glu Glu Glu Ile Arg Glu Ala Phe Arg	
305 310 315 320	

WO 98/40477

PCT/US98/04978

- 54 -

GTT	TTT	GAC	AAG	GAT	GGG	AAC	GGC	TAC	ATC	AGC	GCT	GCT	GAA	TTA	CGT	1008
Val	Phe	Asp	Lys	Asp	Gly	Asn	Gly	Tyr	Ile	Ser	Ala	Ala	Glu	Leu	Arg	
				325					330					335		
CAC	GTC	ATG	ACA	AAC	CTC	GGG	GAG	AAG	TTA	ACA	GAT	GAA	GAA	GTT	GAT	1056
His	Val	Met	Thr	Asn	Leu	Gly	Glu	Lys	Leu	Thr	Asp	Glu	Glu	Val	Asp	
			340					345					350			
GAA	ATG	ATA	AGG	GAA	GCA	GAT	ATC	GAT	GGT	GAT	GGC	CAA	GTA	AAC	TAT	1104
Glu	Met	Ile	Arg	Glu	Ala	Asp	Ile	Asp	Gly	Asp	Gly	Gln	Val	Asn	Tyr	
		355					360					365				
GAA	GAG	TTT	GTA	CAA	ATG	ATG	ACA	GCA	AAG	GGG	GGG	AAG	AGG	CGC	TGG	1152
Glu	Glu	Phe	Val	Gln	Met	Met	Thr	Ala	Lys	Gly	Gly	Lys	Arg	Arg	Trp	
	370					375					380					
AAG	AAA	AAC	TTC	ATT	GCC	GTC	AGC	GCT	GCC	AAC	CGG	TTC	AAG	AAG	ATC	1200
Lys	Lys	Asn	Phe	Ile	Ala	Val	Ser	Ala	Ala	Asn	Arg	Phe	Lys	Lys	Ile	
	385				390					395					400	
TCC	GAG	CTC	ATG	GTG	AGC	AAG	GGC	GAG	GAG	CTG	TTC	ACC	GGG	GTG	GTG	1248
Ser	Glu	Leu	Met	Val	Ser	Lys	Gly	Glu	Glu	Leu	Phe	Thr	Gly	Val	Val	
			405						410					415		
CCC	ATC	CTG	GTC	GAG	CTG	GAC	GGC	GAC	GTA	AAC	GGC	CAC	AAG	TTC	AGC	1296
Pro	Ile	Leu	Val	Glu	Leu	Asp	Gly	Asp	Val	Asn	Gly	His	Lys	Phe	Ser	
			420					425					430			
GTG	TCC	GGC	GAG	GGC	GAG	GGC	GAT	GCC	ACC	TAC	GGC	AAG	CTG	ACC	CTG	1344
Val	Ser	Gly	Glu	Gly	Glu	Gly	Asp	Ala	Thr	Tyr	Gly	Lys	Leu	Thr	Leu	
		435				440						445				
AAG	TTC	ATC	TGC	ACC	ACC	GGC	AAG	CTG	CCC	GTG	CCC	TGG	CCC	ACC	CTC	1392
Lys	Phe	Ile	Cys	Thr	Thr	Gly	Lys	Leu	Pro	Val	Pro	Trp	Pro	Thr	Leu	
	450					455				460						
GTG	ACC	ACC	CTG	ACC	TAC	GGC	GTG	CAG	TGC	TTC	AGC	CGC	TAC	CCC	GAC	1440
Val	Thr	Thr	Leu	Thr	Tyr	Gly	Val	Gln	Cys	Phe	Ser	Arg	Tyr	Pro	Asp	
	465				470			475							480	
CAC	ATG	AAG	CAG	CAC	GAC	TTC	TTC	AAG	TCC	GCC	ATG	CCC	GAA	GGC	TAC	1488
His	Met	Lys	Gln	His	Asp	Phe	Phe	Lys	Ser	Ala	Met	Pro	Glu	Gly	Tyr	
			485					490						495		
GTC	CAG	GAG	CGC	ACC	ATC	TTC	TTC	AAG	GAC	GAC	GGC	AAC	TAC	AAG	ACC	1536
Val	Gln	Glu	Arg	Thr	Ile	Phe	Phe	Lys	Asp	Asp	Gly	Asn	Tyr	Lys	Thr	
			500					505					510			
CGC	GCC	GAG	GTG	AAG	TTC	GAG	GGC	GAC	ACC	CTG	GTG	AAC	CGC	ATC	GAG	1584
Arg	Ala	Glu	Val	Lys	Phe	Glu	Gly	Asp	Thr	Leu	Val	Asn	Arg	Ile	Glu	
		515					520					525				
CTG	AAG	GGC	ATC	GAC	TTC	AAG	GAG	GAC	GGC	AAC	ATC	CTG	GGG	CAC	AAG	1632
Leu	Lys	Gly	Ile	Asp	Phe	Lys	Glu	Asp	Gly	Asn	Ile	Leu	Gly	His	Lys	
	530					535					540					
CTG	GAG	TAC	AAC	TAC	AAC	AGC	CAC	AAC	GTC	TAT	ATC	ATG	GCC	GAC	AAG	1680
Leu	Glu	Tyr	Asn	Tyr	Asn	Ser	His	Asn	Val	Tyr	Ile	Met	Ala	Asp	Lys	
	545				550				555						560	
CAG	AAG	AAC	GGC	ATC	AAG	GTG	AAC	TTC	AAG	ATC	CGC	CAC	AAC	ATC	GAG	1728
Gln	Lys	Asn	Gly	Ile	Lys	Val	Asn	Phe	Lys	Ile	Arg	His	Asn	Ile	Glu	
			565					570						575		

WO 98/40477

PCT/US98/04978

- 55 -

GAC GGC AGC GTG CAG CTC GCC GAC CAC TAC CAG CAG AAC ACC CCC ATC 1776
Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile
580 585 590

GGC GAC GGC CCC GTG CTG CTG CCC GAC AAC CAC TAC CTG AGC ACC CAG 1824
Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln
595 600 605

TCC GCC CTG AGC AAA GAC CCC AAC GAG AAG CGC GAT CAC ATG GTC CTG 1872
Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu
610 615 620

CTG GAG TTC GTG ACC GCC GCC GGG ATC ACT CTC GGC ATG GAC GAG CTG 1920
Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu
625 630 635 640

TAC AAG CCA AAA AAG AAG AGA AAG GTG GAA GAC GCT TAA
1959

Tyr Lys Pro Lys Lys Lys Arg Lys Val Glu Asp Ala.
645 650

(5) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS
(A) LENGTH: 652 AMINO ACIDS
(B) TYPE: AMINO ACID
(C) STRANDEDNESS:
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu
5 10 15

Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly
20 25 30

Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile
35 40 45

Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr
50 55 60

Leu Thr His Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys
65 70 75 80

Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu
85 90 95

Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu
100 105 110

Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly
115 120 125

Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr
130 135 140

Asn Phe Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn
145 150 155 160

WO 98/40477

PCT/US98/04978

- 56 -

Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser
165 170 175

Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly
180 185 190

Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu
195 200 205

Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe
210 215 220

Val Thr Ala Ala Arg Met His Asp Gln Leu Thr Glu Glu Gln Ile Ala
225 230 235 240

Glu Phe Lys Glu Ala Phe Ser Leu Phe Asp Lys Asp Gly Asp Gly Thr
245 250 255

Ile Thr Thr Lys Glu Leu Gly Thr Val Met Arg Ser Leu Gly Gln Asn
260 265 270

Pro Thr Glu Ala Glu Leu Gln Asp Met Ile Asn Glu Val Asp Ala Asp
275 280 285

Gly Asn Gly Thr Ile Tyr Phe Pro Glu Phe Leu Thr Met Met Ala Arg
290 295 300

Lys Met Lys Asp Thr Asp Ser Glu Glu Glu Ile Arg Glu Ala Phe Arg
305 310 315 320

Val Phe Asp Lys Asp Gly Asn Gly Tyr Ile Ser Ala Ala Glu Leu Arg
325 330 335

His Val Met Thr Asn Leu Gly Glu Lys Leu Thr Asp Glu Glu Val Asp
340 345 350

Glu Met Ile Arg Glu Ala Asp Ile Asp Gly Asp Gly Gln Val Asn Tyr
355 360 365

Glu Glu Phe Val Gln Met Met Thr Ala Lys Gly Gly Lys Arg Arg Trp
370 375 380

Lys Lys Asn Phe Ile Ala Val Ser Ala Ala Asn Arg Phe Lys Lys Ile
385 390 395 400

Ser Glu Leu Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val
405 410 415

Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser
420 425 430

Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu
435 440 445

Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu
450 455 460

Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp
465 470 475 480

His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr
485 490 495

WO 98/40477

PCT/US98/04978

- 57 -

Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr
500 505 510

Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu
515 520 525

Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys
530 535 540

Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys
545 550 555 560

Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu
565 570 575

Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile
580 585 590

Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln
595 600 605

Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu
610 615 620

Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu
625 630 635 640

Tyr Lys Pro Lys Lys Lys Arg Lys Val Glu Asp Ala
645 650

(6) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS
(A) LENGTH: 1929 BASE PAIRS
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: SINGLE
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATG GTG AGC AAG GGC GAG GAG CTG TTC ACC GGG GTG GTG CCC ATC CTG	48
Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu	
5 10 15	
GTC GAG CTG GAC GGC GAC GTA AAC GGC CAC AAG TTC AGC GTG TCC GGC	96
Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly	
20 25 30	
GAG GGC GAG GGC GAT GCC ACC TAC GGC AAG CTG ACC CTG AAG TTC ATC	144
Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile	
35 40 45	
TGC ACC ACC GGC AAG CTG CCC GTG CCC TGG CCC ACC CTC GTG ACC ACC	192
Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr	
50 55 60	

WO 98/40477

PCT/US98/04978

- 58 -

CTG	ACC	CAT	GGC	GTG	CAG	TGC	TTC	AGC	CGC	TAC	CCC	GAC	CAC	ATG	AAG	240
Leu	Thr	His	Gly	Val	Gln	Cys	Phe	Ser	Arg	Tyr	Pro	Asp	His	Met	Lys	
65					70					75					80	
CAG	CAC	GAC	TTC	TTC	AAG	TCC	GCC	ATG	CCC	GAA	GGC	TAC	GTC	CAG	GAG	288
Gln	His	Asp	Phe	Phe	Lys	Ser	Ala	Met	Pro	Glu	Gly	Tyr	Val	Gln	Glu	
				85					90					95		
CGC	ACC	ATC	TTC	TTC	AAG	GAC	GAC	GGC	AAC	TAC	AAG	ACC	CGC	GCC	GAG	336
Arg	Thr	Ile	Phe	Phe	Lys	Asp	Asp	Gly	Asn	Tyr	Lys	Thr	Arg	Ala	Glu	
			100					105					110			
GTG	AAG	TTC	GAG	GGC	GAC	ACC	CTG	GTG	AAC	CGC	ATC	GAG	CTG	AAG	GGC	384
Val	Lys	Phe	Glu	Gly	Asp	Thr	Leu	Val	Asn	Arg	Ile	Glu	Leu	Lys	Gly	
		115					120					125				
ATC	GAC	TTC	AAG	GAG	GAC	GGC	AAC	ATC	CTG	GGG	CAC	AAG	CTG	GAG	TAC	432
Ile	Asp	Phe	Lys	Glu	Asp	Gly	Asn	Ile	Leu	Gly	His	Lys	Leu	Glu	Tyr	
	130					135					140					
AAC	TTC	AAC	AGC	CAC	AAC	GTC	TAT	ATC	ATG	GCC	GAC	AAG	CAG	AAG	AAC	480
Asn	Phe	Asn	Ser	His	Asn	Val	Tyr	Ile	Met	Ala	Asp	Lys	Gln	Lys	Asn	
145					150					155					160	
GGC	ATC	AAG	GTG	AAC	TTC	AAG	ATC	CGC	CAC	AAC	ATC	GAG	GAC	GGC	AGC	528
Gly	Ile	Lys	Val	Asn	Phe	Lys	Ile	Arg	His	Asn	Ile	Glu	Asp	Gly	Ser	
				165				170						175		
GTG	CAG	CTC	GCC	GAC	CAC	TAC	CAG	CAG	AAC	ACC	CCC	ATC	GGC	GAC	GGC	576
Val	Gln	Leu	Ala	Asp	His	Tyr	Gln	Gln	Asn	Thr	Pro	Ile	Gly	Asp	Gly	
			180				185						190			
CCC	GTG	CTG	CTG	CCC	GAC	AAC	CAC	TAC	CTG	AGC	ACC	CAG	TCC	GCC	CTG	624
Pro	Val	Leu	Leu	Pro	Asp	Asn	His	Tyr	Leu	Ser	Thr	Gln	Ser	Ala	Leu	
		195				200						205				
AGC	AAA	GAC	CCC	AAC	GAG	AAG	CGC	GAT	CAC	ATG	GTC	CTG	CTG	GAG	TTC	672
Ser	Lys	Asp	Pro	Asn	Glu	Lys	Arg	Asp	His	Met	Val	Leu	Leu	Glu	Phe	
	210					215					220					
GTG	ACC	GCC	GCC	CGC	ATG	CAT	GAC	CAA	CTG	ACA	GAA	GAG	CAG	ATT	GCA	720
Val	Thr	Ala	Ala	Arg	Met	His	Asp	Gln	Leu	Thr	Glu	Glu	Gln	Ile	Ala	
225					230					235					240	
GAG	TTC	AAA	GAA	GCC	TTC	TCA	TTA	TTC	GAC	AAG	GAT	GGG	GAC	GGC	ACC	768
Glu	Phe	Lys	Glu	Ala	Phe	Ser	Leu	Phe	Asp	Lys	Asp	Gly	Asp	Gly	Thr	
				245					250					255		
ATC	ACC	ACA	AAG	GAA	CTT	GGC	ACC	GTT	ATG	AGG	TCG	CTT	GGA	CAA	AAC	816
Ile	Thr	Thr	Lys	Glu	Leu	Gly	Thr	Val	Met	Arg	Ser	Leu	Gly	Gln	Asn	
			260					265					270			
CCA	ACG	GAA	GCA	GAA	TTG	CAG	GAT	ATG	ATC	AAT	GAA	GTC	GAT	GCT	GAT	864
Pro	Thr	Glu	Ala	Glu	Leu	Gln	Asp	Met	Ile	Asn	Glu	Val	Asp	Ala	Asp	
		275					280					285				
GGC	AAT	GGA	ACG	ATT	TAC	TTT	CCT	GAA	TTT	CTT	ACT	ATG	ATG	GCT	AGA	912
Gly	Asn	Gly	Thr	Ile	Tyr	Phe	Pro	Glu	Phe	Leu	Thr	Met	Met	Ala	Arg	
	290					295					300					
AAA	ATG	AAG	GAC	ACA	GAC	AGC	GAA	GAG	GAA	ATC	CGA	GAA	GCA	TTC	CGT	960
Lys	Met	Lys	Asp	Thr	Asp	Ser	Glu	Glu	Glu	Ile	Arg	Glu	Ala	Phe	Arg	
305					310					315					320	

WO 98/40477

PCT/US98/04978

- 59 -

GTT	TTT	GAC	AAG	GAT	GGG	AAC	GGC	TAC	ATC	AGC	GCT	GCT	CAG	TTA	CGT	1008
Val	Phe	Asp	Lys	Asp	Gly	Asn	Gly	Tyr	Ile	Ser	Ala	Ala	Gln	Leu	Arg	
				325					330					335		
CAC	GTC	ATG	ACA	AAC	CTC	GGG	GAG	AAG	TTA	ACA	GAT	GAA	GAA	GTT	GAT	1056
His	Val	Met	Thr	Asn	Leu	Gly	Glu	Lys	Leu	Thr	Asp	Glu	Glu	Val	Asp	
			340					345					350			
GAA	ATG	ATA	AGG	GAA	GCA	GAT	ATC	GAT	GGT	GAT	GGC	CAA	GTA	AAC	TAT	1104
Glu	Met	Ile	Arg	Glu	Ala	Asp	Ile	Asp	Gly	Asp	Gly	Gln	Val	Asn	Tyr	
		355					360					365				
GAA	GAG	TTT	GTA	CAA	ATG	ATG	ACA	GCA	AAG	GGG	GGG	AAG	AGG	CGC	TGG	1152
Glu	Glu	Phe	Val	Gln	Met	Met	Thr	Ala	Lys	Gly	Gly	Lys	Arg	Arg	Trp	
	370					375					380					
AAG	AAA	AAC	TTC	ATT	GCC	GTC	AGC	GCT	GCC	AAC	CGG	TTC	AAG	AAG	ATC	1200
Lys	Lys	Asn	Phe	Ile	Ala	Val	Ser	Ala	Ala	Asn	Arg	Phe	Lys	Lys	Ile	
385					390					395					400	
TCC	GAG	CTC	ATG	GTG	AGC	AAG	GGC	GAG	GAG	CTG	TTC	ACC	GGG	GTG	GTG	1248
Ser	Glu	Leu	Met	Val	Ser	Lys	Gly	Glu	Glu	Leu	Phe	Thr	Gly	Val	Val	
				405					410					415		
CCC	ATC	CTG	GTC	GAG	CTG	GAC	GGC	GAC	GTA	AAC	GGC	CAC	AAG	TTC	AGC	1296
Pro	Ile	Leu	Val	Glu	Leu	Asp	Gly	Asp	Val	Asn	Gly	His	Lys	Phe	Ser	
			420					425					430			
GTG	TCC	GGC	GAG	GGC	GAG	GGC	GAT	GCC	ACC	TAC	GGC	AAG	CTG	ACC	CTG	1344
Val	Ser	Gly	Glu	Gly	Glu	Gly	Asp	Ala	Thr	Tyr	Gly	Lys	Leu	Thr	Leu	
		435					440					445				
AAG	TTC	ATC	TGC	ACC	ACC	GGC	AAG	CTG	CCC	GTG	CCC	TGG	CCC	ACC	CTC	1392
Lys	Phe	Ile	Cys	Thr	Thr	Gly	Lys	Leu	Pro	Val	Pro	Trp	Pro	Thr	Leu	
	450					455					460					
GTG	ACC	ACC	CTG	ACC	TAC	GGC	GTG	CAG	TGC	TTC	AGC	CGC	TAC	CCC	GAC	1440
Val	Thr	Thr	Leu	Thr	Tyr	Gly	Val	Gln	Cys	Phe	Ser	Arg	Tyr	Pro	Asp	
465					470				475						480	
CAC	ATG	AAG	CAG	CAC	GAC	TTC	TTC	AAG	TCC	GCC	ATG	CCC	GAA	GGC	TAC	1488
His	Met	Lys	Gln	His	Asp	Phe	Phe	Lys	Ser	Ala	Met	Pro	Glu	Gly	Tyr	
				485					490					495		
GTC	CAG	GAG	CGC	ACC	ATC	TTC	TTC	AAG	GAC	GAC	GGC	AAC	TAC	AAG	ACC	1536
Val	Gln	Glu	Arg	Thr	Ile	Phe	Phe	Lys	Asp	Asp	Gly	Asn	Tyr	Lys	Thr	
			500					505					510			
CGC	GCC	GAG	GTG	AAG	TTC	GAG	GGC	GAC	ACC	CTG	GTG	AAC	CGC	ATC	GAG	1584
Arg	Ala	Glu	Val	Lys	Phe	Glu	Gly	Asp	Thr	Leu	Val	Asn	Arg	Ile	Glu	
		515					520					525				
CTG	AAG	GGC	ATC	GAC	TTC	AAG	GAG	GAC	GGC	AAC	ATC	CTG	GGG	CAC	AAG	1632
Leu	Lys	Gly	Ile	Asp	Phe	Lys	Glu	Asp	Gly	Asn	Ile	Leu	Gly	His	Lys	
	530					535					540					
CTG	GAG	TAC	AAC	TAC	AAC	AGC	CAC	AAC	GTC	TAT	ATC	ATG	GCC	GAC	AAG	1680
Leu	Glu	Tyr	Asn	Tyr	Asn	Ser	His	Asn	Val	Tyr	Ile	Met	Ala	Asp	Lys	
545					550				555						560	
CAG	AAG	AAC	GGC	ATC	AAG	GTG	AAC	TTC	AAG	ATC	CGC	CAC	AAC	ATC	GAG	1728
Gln	Lys	Asn	Gly	Ile	Lys	Val	Asn	Phe	Lys	Ile	Arg	His	Asn	Ile	Glu	
				565					570					575		

WO 98/40477

PCT/US98/04978

- 60 -

GAC	GGC	AGC	GTG	CAG	CTC	GCC	GAC	CAC	TAC	CAG	CAG	AAC	ACC	CCC	ATC	1776	
Asp	Gly	Ser	Val	Gln	Leu	Ala	Asp	His	Tyr	Gln	Gln	Asn	Thr	Pro	Ile		
			580					585					590				
GGC	GAC	GGC	CCC	GTG	CTG	CTG	CCC	GAC	AAC	CAC	TAC	CTG	AGC	ACC	CAG	1824	
Gly	Asp	Gly	Pro	Val	Leu	Leu	Pro	Asp	Asn	His	Tyr	Leu	Ser	Thr	Gln		
		595					600					605					
TCC	GCC	CTG	AGC	AAA	GAC	CCC	AAC	GAG	AAG	CGC	GAT	CAC	ATG	GTC	CTG	1872	
Ser	Ala	Leu	Ser	Lys	Asp	Pro	Asn	Glu	Lys	Arg	Asp	His	Met	Val	Leu		
	610					615					620						
CTG	GAG	TTC	GTG	ACC	GCC	GCC	GGG	ATC	ACT	CTC	GGC	ATG	GAC	GAG	CTG	1920	
Leu	Glu	Phe	Val	Thr	Ala	Ala	Gly	Ile	Thr	Leu	Gly	Met	Asp	Glu	Leu		
625					630					635					640		
TAC	AAG	TAA															
	1929																
Tyr	Lys																

(7) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS
 (A) LENGTH: 642 AMINO ACIDS
 (B) TYPE: AMINO ACID
 (C) STRANDEDNESS:
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met	Val	Ser	Lys	Gly	Glu	Glu	Leu	Phe	Thr	Gly	Val	Val	Pro	Ile	Leu	
				5					10					15		
Val	Glu	Leu	Asp	Gly	Asp	Val	Asn	Gly	His	Lys	Phe	Ser	Val	Ser	Gly	
			20					25					30			
Glu	Gly	Glu	Gly	Asp	Ala	Thr	Tyr	Gly	Lys	Leu	Thr	Leu	Lys	Phe	Ile	
		35					40					45				
Cys	Thr	Thr	Gly	Lys	Leu	Pro	Val	Pro	Trp	Pro	Thr	Leu	Val	Thr	Thr	
	50				55					60						
Leu	Thr	His	Gly	Val	Gln	Cys	Phe	Ser	Arg	Tyr	Pro	Asp	His	Met	Lys	
	65				70				75					80		
Gln	His	Asp	Phe	Phe	Lys	Ser	Ala	Met	Pro	Glu	Gly	Tyr	Val	Gln	Glu	
			85					90						95		
Arg	Thr	Ile	Phe	Phe	Lys	Asp	Asp	Gly	Asn	Tyr	Lys	Thr	Arg	Ala	Glu	
			100				105						110			
Val	Lys	Phe	Glu	Gly	Asp	Thr	Leu	Val	Asn	Arg	Ile	Glu	Leu	Lys	Gly	
		115				120						125				
Ile	Asp	Phe	Lys	Glu	Asp	Gly	Asn	Ile	Leu	Gly	His	Lys	Leu	Glu	Tyr	
	130				135						140					
Asn	Phe	Asn	Ser	His	Asn	Val	Tyr	Ile	Met	Ala	Asp	Lys	Gln	Lys	Asn	
	145				150				155						160	
Gly	Ile	Lys	Val	Asn	Phe	Lys	Ile	Arg	His	Asn	Ile	Glu	Asp	Gly	Ser	
			165					170						175		

WO 98/40477

PCT/US98/04978

- 61 -

Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly
180 185 190

Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu
195 200 205

Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe
210 215 220

Val Thr Ala Ala Arg Met His Asp Gln Leu Thr Glu Glu Gln Ile Ala
225 230 235 240

Glu Phe Lys Glu Ala Phe Ser Leu Phe Asp Lys Asp Gly Asp Gly Thr
245 250 255

Ile Thr Thr Lys Glu Leu Gly Thr Val Met Arg Ser Leu Gly Gln Asn
260 265 270

Pro Thr Glu Ala Glu Leu Gln Asp Met Ile Asn Glu Val Asp Ala Asp
275 280 285

Gly Asn Gly Thr Ile Tyr Phe Pro Glu Phe Leu Thr Met Met Ala Arg
290 295 300

Lys Met Lys Asp Thr Asp Ser Glu Glu Glu Ile Arg Glu Ala Phe Arg
305 310 315 320

Val Phe Asp Lys Asp Gly Asn Gly Tyr Ile Ser Ala Ala Gln Leu Arg
325 330 335

His Val Met Thr Asn Leu Gly Glu Lys Leu Thr Asp Glu Glu Val Asp
340 345 350

Glu Met Ile Arg Glu Ala Asp Ile Asp Gly Asp Gly Gln Val Asn Tyr
355 360 365

Glu Glu Phe Val Gln Met Met Thr Ala Lys Gly Gly Lys Arg Arg Trp
370 375 380

Lys Lys Asn Phe Ile Ala Val Ser Ala Ala Asn Arg Phe Lys Lys Ile
385 390 395 400

Ser Glu Leu Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val
405 410 415

Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser
420 425 430

Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu
435 440 445

Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu
450 455 460

Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp
465 470 475 480

His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr
485 490 495

Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr
500 505 510

Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu
515 520 525

WO 98/40477

PCT/US98/04978

- 62 -

Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys
 530 535 540
 Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys
 545 550 555 560
 Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu
 565 570 575
 Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile
 580 585 590
 Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln
 595 600 605
 Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu
 610 615 620
 Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu
 625 630 635 640
 Tyr Lys

(8) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS
- (A) LENGTH: 1971 BASE PAIRS
 - (B) TYPE: NUCLEIC ACID
 - (C) STRANDEDNESS: SINGLE
 - (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ATG CTG CTG CCC GTC CCC CTG CTG CTG GGC CTG CTG GGC GCC GCC GCC	48
Met Leu Leu Pro Val Pro Leu Leu Leu Gly Leu Leu Gly Ala Ala Ala	
5 10 15	
GAC GTG AGC AAG GGC GAG GAG CTG TTC ACC GGG GTG GTG CCC ATC CTG	96
Asp Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu	
20 25 30	
GTC GAG CTG GAC GGC GAC GTA AAC GGC CAC AAG TTC AGC GTG TCC GGC	144
Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly	
35 40 45	
GAG GGC GAG GGC GAT GCC ACC TAC GGC AAG CTG ACC CTG AAG TTC ATC	192
Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile	
50 55 60	
TGC ACC ACC GGC AAG CTG CCC GTG CCC TGG CCC ACC CTC GTG ACC ACC	240
Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr	
65 70 75 80	
CTG ACC CAT GGC GTG CAG TGC TTC AGC CGC TAC CCC GAC CAC ATG AAG	288
Leu Thr His Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys	
85 90 95	
CAG CAC GAC TTC TTC AAG TCC GCC ATG CCC GAA GGC TAC GTC CAG GAG	336
Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu	
100 105 110	

WO 98/40477

PCT/US98/04978

- 63 -

CGC ACC ATC TTC TTC AAG GAC GAC GGC AAC TAC AAG ACC CGC GCC GAG	384
Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu	
115 120 125	
GTG AAG TTC GAG GGC GAC ACC CTG GTG AAC CGC ATC GAG CTG AAG GGC	432
Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly	
130 135 140	
ATC GAC TTC AAG GAG GAC GGC AAC ATC CTG GGG CAC AAG CTG GAG TAC	480
Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr	
145 150 155 160	
AAC TTC AAC AGC CAC AAC GTC TAT ATC ATG GCC GAC AAG CAG AAG AAC	528
Asn Phe Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn	
165 170 175	
GGC ATC AAG GTG AAC TTC AAG ATC CGC CAC AAC ATC GAG GAC GGC AGC	576
Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser	
180 185 190	
GTG CAG CTC GCC GAC CAC TAC CAG CAG AAC ACC CCC ATC GGC GAC GGC	624
Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly	
195 200 205	
CCC GTG CTG CTG CCC GAC AAC CAC TAC CTG AGC ACC CAG TCC GCC CTG	672
Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu	
210 215 220	
AGC AAA GAC CCC AAC GAG AAG CGC GAT CAC ATG GTC CTG CTG GAG TTC	720
Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe	
225 230 235 240	
GTG ACC GCC GCC CGC ATG CAT GAC CAA CTG ACA GAA GAG CAG ATT GCA	768
Val Thr Ala Ala Arg Met His Asp Gln Leu Thr Glu Glu Gln Ile Ala	
245 250 255	
GAG TTC AAA GAA GCC TTC TCA TTA TTC GAC AAG GAT GGG GAC GGC ACC	816
Glu Phe Lys Glu Ala Phe Ser Leu Phe Asp Lys Asp Gly Asp Gly Thr	
260 265 270	
ATC ACC ACA AAG GAA CTT GGC ACC GTT ATG AGG TCG CTT GGA CAA AAC	864
Ile Thr Thr Lys Glu Leu Gly Thr Val Met Arg Ser Leu Gly Gln Asn	
275 280 285	
CCA ACG GAA GCA GAA TTG CAG GAT ATG ATC AAT GAA GTC GAT GCT GAT	912
Pro Thr Glu Ala Glu Leu Gln Asp Met Ile Asn Glu Val Asp Ala Asp	
290 295 300	
GGC AAT GGA ACG ATT TAC TTT CCT GAA TTT CTT ACT ATG ATG GCT AGA	960
Gly Asn Gly Thr Ile Tyr Phe Pro Glu Phe Leu Thr Met Met Ala Arg	
305 310 315 320	
AAA ATG AAG GAC ACA GAC AGC GAA GAG GAA ATC CGA GAA GCA TTC CGT	1008
Lys Met Lys Asp Thr Asp Ser Glu Glu Glu Ile Arg Glu Ala Phe Arg	
325 330 335	
GTT TTT GAC AAG GAT GGG AAC GGC TAC ATC AGC GCT GCT CAG TTA CGT	1056
Val Phe Asp Lys Asp Gly Asn Gly Tyr Ile Ser Ala Ala Gln Leu Arg	
340 345 350	
CAC GTC ATG ACA AAC CTC GGG GAG AAG TTA ACA GAT GAA GAA GTT GAT	1104
His Val Met Thr Asn Leu Gly Glu Lys Leu Thr Asp Glu Glu Val Asp	
355 360 365	

WO 98/40477

PCT/US98/04978

- 64 -

GAA Glu 370	ATG Met	ATA Ile	AGG Arg	GAA Glu	GCA Ala	GAT Asp 375	ATC Ile	GAT Asp	GGT Gly	GAT Asp	GGC Gly 380	CAA Gln	GTA Val	AAC Asn	TAT Tyr	1152
GAA Glu 385	GAG Glu	TTT Phe	GTA Val	CAA Gln	ATG Met 390	ATG Met	ACA Thr	GCA Ala	AAG Lys	GGG Gly 395	GGG Gly	AAG Lys	AGG Arg	CGC Arg	TGG Trp 400	1200
AAG Lys	AAA Lys	AAC Asn	TTC Phe	ATT Ile 405	GCC Ala	GTC Val	AGC Ser	GCT Ala	GCC Ala 410	AAC Asn	CGG Arg	TTC Phe	AAG Lys 415	AAG Lys	ATC Ile	1248
TCC Ser	GAG Glu	CTC Leu	ATG Met 420	GTG Val	AGC Ser	AAG Lys	GGC Gly	GAG Glu	GAG Glu	CTG Leu	TTC Phe	ACC Thr	GGG Gly 430	GTG Val	GTG Val	1296
CCC Pro	ATC Ile	CTG Leu 435	GTC Val	GAG Glu	CTG Leu	GAC Asp	GGC Gly 440	GAC Asp	GTA Val	AAC Asn	GGC Gly	CAC His 445	AAG Lys	TTC Phe	AGC Ser	1344
GTG Val 450	TCC Ser	GGC Gly	GAG Glu	GGC Gly	GAG Glu	GGC Gly 455	GAT Asp	GCC Ala	ACC Thr	TAC Tyr	GGC Gly 460	AAG Lys	CTG Leu	ACC Thr	CTG Leu	1392
AAG Lys 465	TTC Phe	ATC Ile	TGC Cys	ACC Thr	ACC Thr 470	GGC Gly	AAG Lys	CTG Leu	CCC Pro	GTG Val 475	CCC Pro	TGG Trp	CCC Pro	ACC Thr	CTC Leu 480	1440
GTG Val	ACC Thr	ACC Thr	CTG Leu	ACC Thr	TAC Tyr 485	GGC Gly	GTG Val	CAG Gln	TGC Cys 490	TTC Phe	AGC Ser	CGC Arg	TAC Tyr	CCC Pro 495	GAC Asp	1488
CAC His	ATG Met	AAG Lys	CAG Gln 500	CAC His	GAC Asp	TTC Phe	TTC Phe 505	AAG Lys	TCC Ser	GCC Ala	ATG Met	CCC Pro 510	GAA Glu	GGC Gly	TAC Tyr	1536
GTC Val	CAG Gln 515	GAG Glu	CGC Arg	ACC Thr	ATC Ile	TTC Phe 520	TTC Phe	AAG Lys	GAC Asp	GAC Asp	GGC Gly 525	AAC Asn	TAC Tyr	AAG Lys	ACC Thr	1584
CGC Arg 530	GCC Ala	GAG Glu	GTG Val	AAG Lys	TTC Phe	GAG Glu 535	GGC Gly	GAC Asp	ACC Thr	CTG Leu	GTG Val 540	AAC Asn	CGC Arg	ATC Ile	GAG Glu	1632
CTG Leu 545	AAG Lys	GGC Gly	ATC Ile	GAC Asp	TTC Phe 550	AAG Lys	GAG Glu	GAC Asp	GGC Gly 555	AAC Asn	ATC Ile	CTG Leu	GGG Gly	CAC His	AAG Lys 560	1680
CTG Leu	GAG Glu	TAC Tyr	AAC Asn 565	TAC Tyr	AAC Asn	AGC Ser	CAC His	AAC Asn	GTC Val 570	TAT Tyr	ATC Ile	ATG Met	GCC Ala 575	GAC Asp	AAG Lys	1728
CAG Gln	AAG Lys	AAC Asn	GGC Gly 580	ATC Ile	AAG Lys	GTG Val	AAC Asn 585	TTC Phe	AAG Lys	ATC Ile	CGC Arg	CAC His	AAC Asn 590	ATC Ile	GAG Glu	1776
GAC Asp	GGC Gly 595	AGC Ser	GTG Val	CAG Gln	CTC Leu	GCC Ala 600	GAC Asp	CAC His	TAC Tyr	CAG Gln	CAG Gln	AAC Asn 605	ACC Thr	CCC Pro	ATC Ile	1824
GGC Gly 610	GAC Asp	GGC Gly	CCC Pro	GTG Val	CTG Leu	CTG Leu 615	CCC Pro	GAC Asp	AAC Asn	CAC His	TAC Tyr 620	CTG Leu	AGC Ser	ACC Thr	CAG Gln	1872

WO 98/40477

PCT/US98/04978

- 65 -

TCC GCC CTG AGC AAA GAC CCC AAC GAG AAG CGC GAT CAC ATG GTC CTG 1920
 Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu
 625 630 635 640

CTG GAG TTC GTG ACC GCC GCC GGG ATC ACT CTC GGC AAG GAC GAG CTG 1968
 Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu Gly Lys Asp Glu Leu
 645 650 655

TAA 1971

(9) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS
 (A) LENGTH: 656 AMINO ACIDS
 (B) TYPE: AMINO ACID
 (C) STRANDEDNESS:
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Leu Leu Pro Val Pro Leu Leu Leu Gly Leu Leu Gly Ala Ala Ala
 5 10 15

Asp Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu
 20 25 30

Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly
 35 40 45

Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile
 50 55 60

Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr
 65 70 75 80

Leu Thr His Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys
 85 90 95

Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu
 100 105 110

Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu
 115 120 125

Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly
 130 135 140

Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr
 145 150 155 160

Asn Phe Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn
 165 170 175

Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser
 180 185 190

Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly
 195 200 205

Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu
 210 215 220

WO 98/40477

PCT/US98/04978

- 66 -

Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe
225 230 235 240

Val Thr Ala Ala Arg Met His Asp Gln Leu Thr Glu Glu Gln Ile Ala
245 250 255

Glu Phe Lys Glu Ala Phe Ser Leu Phe Asp Lys Asp Gly Asp Gly Thr
260 265 270

Ile Thr Thr Lys Glu Leu Gly Thr Val Met Arg Ser Leu Gly Gln Asn
275 280 285

Pro Thr Glu Ala Glu Leu Gln Asp Met Ile Asn Glu Val Asp Ala Asp
290 295 300

Gly Asn Gly Thr Ile Tyr Phe Pro Glu Phe Leu Thr Met Met Ala Arg
305 310 315 320

Lys Met Lys Asp Thr Asp Ser Glu Glu Glu Ile Arg Glu Ala Phe Arg
325 330 335

Val Phe Asp Lys Asp Gly Asn Gly Tyr Ile Ser Ala Ala Gln Leu Arg
340 345 350

His Val Met Thr Asn Leu Gly Glu Lys Leu Thr Asp Glu Glu Val Asp
355 360 365

Glu Met Ile Arg Glu Ala Asp Ile Asp Gly Asp Gly Gln Val Asn Tyr
370 375 380

Glu Glu Phe Val Gln Met Met Thr Ala Lys Gly Gly Lys Arg Arg Trp
385 390 395 400

Lys Lys Asn Phe Ile Ala Val Ser Ala Ala Asn Arg Phe Lys Lys Ile
405 410 415

Ser Glu Leu Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val
420 425 430

Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser
435 440 445

Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu
450 455 460

Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu
465 470 475 480

Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp
485 490 495

His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr
500 505 510

Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr
515 520 525

Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu
530 535 540

Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys
545 550 555 560

WO 98/40477

PCT/US98/04978

- 67 -

Leu	Glu	Tyr	Asn	Tyr	Asn	Ser	His	Asn	Val	Tyr	Ile	Met	Ala	Asp	Lys
				565					570					575	
Gln	Lys	Asn	Gly	Ile	Lys	Val	Asn	Phe	Lys	Ile	Arg	His	Asn	Ile	Glu
			580					585					590		
Asp	Gly	Ser	Val	Gln	Leu	Ala	Asp	His	Tyr	Gln	Gln	Asn	Thr	Pro	Ile
		595					600					605			
Gly	Asp	Gly	Pro	Val	Leu	Leu	Pro	Asp	Asn	His	Tyr	Leu	Ser	Thr	Gln
	610					615					620				
Ser	Ala	Leu	Ser	Lys	Asp	Pro	Asn	Glu	Lys	Arg	Asp	His	Met	Val	Leu
625					630					635					640
Leu	Glu	Phe	Val	Thr	Ala	Ala	Gly	Ile	Thr	Leu	Gly	Lys	Asp	Glu	Leu
				645					650					655	

WO 98/40477**PCT/US98/04978**

- 68 -

From the above description, the essential characteristics of the present invention can be ascertained. Without departing from the spirit and scope thereof, various changes and modifications of the invention can be made to adapt it to various usages and conditions. Thus, other embodiments are also within the claims.

WO 98/40477

PCT/US98/04978

- 69 -

What is claimed is:

1. A nucleic acid molecule, comprising a polynucleotide that encodes a fluorescent protein energy transfer partner, comprising:
 - a first chemical entity binding region that binds to a chemical entity and binds to a second protein partner, and
 - a fluorescent protein in frame with said first chemical entity binding region; wherein said fluorescent protein is a first energy transfer partner for a second energy transfer partner attached to said second protein partner.
2. The nucleic acid molecule of claim 1, wherein said fluorescent protein is an *Aequorea*-related fluorescent protein.
3. The nucleic acid molecule of claim 1, wherein said first chemical entity binding region comprises a bound conformation of smaller volume when said chemical entity is bound to said first chemical entity binding region compared to said first chemical entity binding region when said chemical entity is not bound to said first chemical entity binding region and said bound conformation permits increased energy transfer between said first energy transfer partner and said second energy transfer partner.
4. The nucleic acid molecule of claim 3, wherein said bound conformation increases the affinity between said first chemical entity binding region and said second protein partner.
5. The nucleic acid molecule of claim 2, wherein said first chemical entity binding region binds to a protein association region of said second protein partner.
6. The nucleic acid molecule of claim 3, wherein said chemical entity is an analyte.

WO 98/40477

PCT/US98/04978

- 70 -

7. The nucleic acid molecule of claim 2, wherein said first chemical entity binding region is either calmodulin, a calmodulin-related protein moiety, cGMP-dependent protein kinase, a steroid hormone receptor, a ligand binding domain of a steroid hormone receptor, protein kinase C, inositol-1,4,5-triphosphate receptor, or recoverin.
8. The nucleic acid molecule of claim 3, wherein said first chemical entity binding region is calmodulin or a calmodulin-related protein.
9. The nucleic acid molecule of claim 1, wherein said fluorescent protein energy transfer partner further comprises a second energy transfer partner in frame with said first energy transfer partner.
10. The nucleic acid molecule of claim 9, wherein said first energy transfer partner is either P4-3, EBFP, or W1B, and said second energy transfer partner is either S65T, EGFP, or 10c.
11. The nucleic acid molecule of claim 1, wherein said fluorescent protein energy transfer partner further comprises a second protein partner comprises said second energy transfer partner in frame with said first energy transfer partner.
12. The nucleic acid molecule of claim 11, wherein said second protein partner further comprises a protein association region.
13. The nucleic acid molecule of claim 11, wherein said second protein partner is either a calmodulin-binding domain of skMLCKp, smMLCK, CaMKII, Caldesmon, Calspermin, phosphofructokinase calcineurin, phosphorylase kinase, Ca²⁺-ATPase 59 kDa PDE, 60 kDa PDE, nitric oxide synthase, type I adenylyl cyclase, *Bordetella pertussis* adenylyl cyclase, Neuromodulin, Spectrin, MARCKS, F52, β -Adducin, HSP90a, HIV-1 gp160, BBMHBI, Dilute MHC, Mastoparan, Melittin, Glucagon, Secretin, VIP, GIP, or Model Peptide CBP2.

WO 98/40477

PCT/US98/04978

- 71 -

14. The nucleic acid molecule of claim 13, wherein the target peptide moiety is M13.
15. The nucleic acid molecule of claim 11, further comprising an amino acid linker in frame with said first chemical entity binding region and fusing said protein association region with said first chemical entity binding region.
16. The nucleic acid of claim 15, wherein said fluorescent protein energy transfer partner further comprises a localization sequence.
17. A nucleic acid molecule, comprising a polynucleotide that encodes:
 - 1) a first fluorescent protein energy transfer partner, comprising:
 - a chemical entity binding region that binds to a chemical entity and binds to a second fluorescent protein energy transfer partner, and
 - a first fluorescent protein in frame with said first chemical entity binding region; and
 - 2) said second fluorescent protein energy transfer partner, comprising:
 - a protein association region that binds to said chemical entity binding region, and
 - a second fluorescent protein in frame with said protein association region;wherein said first fluorescent protein and said second fluorescent protein are energy transfer partners and said first fluorescent protein energy transfer partner is in frame with said second first fluorescent protein energy transfer partner.
18. The nucleic acid molecule of claim 17, wherein said first and fluorescent protein are *Aequorea*-related fluorescent proteins.

WO 98/40477

PCT/US98/04978

- 72 -

19. The nucleic acid molecule of claim 17, wherein said chemical entity binding region comprises a bound conformation of smaller volume when said chemical entity is bound to said chemical entity binding region compared to said chemical entity binding region when said chemical entity is not bound to said chemical entity binding region and said bound conformation permits increased energy transfer between said first fluorescent protein energy transfer partner and said second fluorescent protein energy transfer partner.
20. The nucleic acid molecule of claim 19, wherein said bound conformation increases the affinity between said chemical entity binding region and said second fluorescent protein energy transfer partner.
21. The nucleic acid molecule of claim 20, wherein said chemical entity is an analyte.
22. A system for monitoring protein-protein association, comprising:
 - 1) a first fluorescent protein energy transfer partner, comprising:
 - a chemical entity binding region that binds to a chemical entity and binds to a second fluorescent protein energy transfer partner, and
 - a first energy transfer partner; and
 - 2) said second fluorescent protein energy transfer partner, comprising:
 - a protein association region that binds to said chemical entity binding region,and
 - a second energy transfer partner;
 - wherein said first energy transfer partner and said energy transfer partner are energy transfer partners.
23. An expression vector containing the nucleic acid sequence of claim 1.
24. A transgenic non-human animal comprising a nucleic acid sequence according to claim 1.

WO 98/40477

PCT/US98/04978

- 73 -

25. An expression vector comprising expression control sequences operatively linked to a nucleic acid sequence coding for the expression of a fluorescent indicator, the indicator comprising:
- a binding protein moiety having an analyte-binding region which binds an analyte and causes the indicator to change conformation upon exposure to the analyte;
 - a donor fluorescent protein moiety fused to the binding protein moiety; and
 - an acceptor fluorescent protein moiety fused to the binding protein moiety, wherein the donor moiety and the acceptor moiety change position relative to each other when the analyte binds to the analyte-binding region, altering fluorescence resonance energy transfer between the donor moiety and the acceptor moiety when the donor moiety is excited.
26. The expression vector of claim 14 adapted for function in a prokaryotic cell.
27. The expression vector of claim 14 adapted for function in a eukaryotic cell.
28. A host cell transfected with an expression vector comprising an expression control sequence operatively linked to a sequence coding for the expression of a fluorescent indicator, the indicator comprising:
- 1. a binding protein moiety having an analyte-binding region which binds an analyte and causes the indicator to change conformation upon exposure to the analyte;
 - 2. a donor fluorescent protein moiety covalently coupled to the binding protein moiety; and
 - 3. an acceptor fluorescent protein moiety covalently coupled to the binding protein moiety,
- wherein the donor moiety and the acceptor moiety change position relative to each other when the analyte binds to the analyte-binding region, altering fluorescence resonance energy transfer between the donor moiety and the acceptor moiety when the donor moiety is excited.

WO 98/40477

PCT/US98/04978

- 74 -

29. The cell of claim 17, wherein the cell is a prokaryote.
30. The cell of claim 18, wherein the cell is *E. coli*.
31. The cell of claim 17, wherein the cell is a eukaryotic cell.
32. The cell of claim 20, wherein the cell is a yeast cell.
33. The cell of claim 20, wherein the cell is a mammalian cell.
34. A transgenic non-human animal having a phenotype characterized by expression of the nucleic acid sequence of claim 1, the phenotype being conferred by a transgene contained in the somatic and germ cells of the mouse, the transgene comprising a nucleic acid sequence which encodes a fluorescent indicator specific antigen polypeptide.
35. The transgenic non-human animal of claim 23, wherein the animal is a mouse.
36. A method for producing a transgenic non-human animal having a phenotype characterized by expression of the nucleic acid sequence of claim 1, the method comprising:
 - (a) introducing a transgene into a zygote of an animal, the transgene comprising a DNA construct encoding a the fluorescent indicator specific antigen;
 - (b) transplanting the zygote into a pseudopregnant animal;
 - (c) allowing the zygote to develop to term; and
 - (d) identifying at least one transgenic offspring containing the transgene.

WO 98/40477

PCT/US98/04978

- 75 -

37. The method of claim 25, wherein the introducing of the transgene into the embryo is by introducing an embryonic stem cell containing the transgene into the embryo.
38. The method of claim 25, wherein the introducing of the transgene into the embryo is by infecting the embryo with a retrovirus containing the transgene.
39. The method of claim 25, wherein the animal is a mouse.
40. An isolated nucleic acid sequence which encodes a fluorescent indicator, the indicator comprising:
- a binding protein moiety having an analyte-binding region which binds an analyte and causes the indicator to change conformation upon exposure to the analyte;
 - a donor fluorescent protein moiety covalently coupled to the binding protein moiety; and
 - an acceptor fluorescent protein moiety covalently coupled to the binding protein moiety,
- wherein the donor moiety and the acceptor moiety change position relative to each other when the analyte binds to the analyte-binding region, altering fluorescence resonance energy transfer between the donor moiety and the acceptor moiety when the donor moiety is excited.
41. The nucleic acid of claim 40, wherein the donor fluorescent protein moiety and the acceptor fluorescent protein moiety are *Aequorea*-related fluorescent protein moieties.
42. The nucleic acid of claim 41, wherein the indicator further includes a target peptide moiety and a linker moiety that covalently couples the binding protein and the target peptide moiety and is a peptide moiety, and the binding protein moiety further includes a peptide-binding region for binding a target peptide moiety.

WO 98/40477

PCT/US98/04978

- 76 -

43. The nucleic acid of claim 42, wherein the donor fluorescent protein moiety is fused to the binding protein moiety and the acceptor fluorescent protein moiety is fused to the target peptide moiety.
44. The nucleic acid of claim 41, wherein the binding protein moiety is calmodulin, a calmodulin-related protein moiety, cGMP-dependent protein kinase, a steroid hormone receptor, a ligand binding domain of a steroid hormone receptor, protein kinase C, inositol-1,4,5-triphosphate receptor, or recoverin.
45. The nucleic acid of claim 42, wherein the binding protein moiety is calmodulin or a calmodulin-related protein moiety.
46. The nucleic acid of claim 45, wherein the donor fluorescent protein moiety is P4-3, EBFP, or W1B, and the acceptor fluorescent protein moiety is S65T, EGFP, or 10c.
47. The nucleic acid of claim 45, wherein the target peptide moiety is a calmodulin-binding domain of skMLCKp, smMLCK, CaMKII, Caldesmon, Calspermin, p-hosphofructokinase calcineurin, phosphorylase kinase, Ca²⁺-ATPase 59 kDa PDE, 60 kDa PDE, nitric oxide synthase, type I adenylyl cyclase, *Bordetella pertussis* adenylyl cyclase, Neuromodulin, Spectrin, MARCKS, F52, β -Adducin, HSP90a, HIV-1 gp160, BBMHB1, Dilute MHC, Mastoparan, Melittin, Glucagon, Secretin, VIP, GIP, or Model Peptide CBP2.
48. The nucleic acid of claim 47, wherein the target peptide moiety is M13.
49. The nucleic acid of claim 48, wherein the linker moiety is -Gly-Gly-.
50. The nucleic acid of claim 40, wherein the indicator further comprises a localization sequence.

WO 98/40477

PCT/US98/04978

- 77 -

51. An expression vector containing the nucleic acid sequence of claim 40.
52. A transgenic non-human animal comprising a nucleic acid sequence according to claim 40.
53. An expression vector comprising expression control sequences operatively linked to a nucleic acid sequence coding for the expression of a fluorescent indicator, the indicator comprising:
 - a binding protein moiety having an analyte-binding region which binds an analyte and causes the indicator to change conformation upon exposure to the analyte;
 - a donor fluorescent protein moiety fused to the binding protein moiety;
 - and
 - an acceptor fluorescent protein moiety fused to the binding protein moiety,
 - wherein the donor moiety and the acceptor moiety change position relative to each other when the analyte binds to the analyte-binding region, altering fluorescence resonance energy transfer between the donor moiety and the acceptor moiety when the donor moiety is excited.
54. The expression vector of claim 53 adapted for function in a prokaryotic cell.
55. The expression vector of claim 53 adapted for function in a eukaryotic cell.

WO 98/40477

PCT/US98/04978

- 78 -

56. A host cell transfected with an expression vector comprising an expression control sequence operatively linked to a sequence coding for the expression of a fluorescent indicator, the indicator comprising:

a binding protein moiety having an analyte-binding region which binds an analyte and causes the indicator to change conformation upon exposure to the analyte;

a donor fluorescent protein moiety covalently coupled to the binding protein moiety; and

an acceptor fluorescent protein moiety covalently coupled to the binding protein moiety,

wherein the donor moiety and the acceptor moiety change position relative to each other when the analyte binds to the analyte-binding region, altering fluorescence resonance energy transfer between the donor moiety and the acceptor moiety when the donor moiety is excited.

57. The cell of claim 56, wherein the cell is a prokaryote.

58. The cell of claim 57, wherein the cell is *E. coli*.

59. The cell of claim 56, wherein the cell is a eukaryotic cell.

60. The cell of claim 59, wherein the cell is a yeast cell.

61. The cell of claim 59, wherein the cell is a mammalian cell.

62. A transgenic non-human animal having a phenotype characterized by expression of the nucleic acid sequence of claim 40, the phenotype being conferred by a transgene contained in the somatic and germ cells of the mouse, the transgene comprising a nucleic acid sequence which encodes a fluorescent indicator specific antigen polypeptide.

WO 98/40477

PCT/US98/04978

- 79 -

63. The transgenic non-human animal of claim 62, wherein the animal is a mouse.
64. A method for producing a transgenic non-human animal having a phenotype characterized by expression of the nucleic acid sequence of claim 40, the method comprising:
- (a) introducing a transgene into a zygote of an animal, the transgene comprising a DNA construct encoding a the fluorescent indicator specific antigen;
 - (b) transplanting the zygote into a pseudopregnant animal;
 - (c) allowing the zygote to develop to term; and
 - (d) identifying at least one transgenic offspring containing the transgene.
65. The method of claim 64, wherein the introducing of the transgene into the embryo is by introducing an embryonic stem cell containing the transgene into the embryo.
66. The method of claim 64, wherein the introducing of the transgene into the embryo is by infecting the embryo with a retrovirus containing the transgene.
67. The method of claim 64, wherein the animal is a mouse.

WO 98/40477

PCT/US98/04978

- 80 -

68. A fluorescent indicator comprising:

a binding protein moiety having an analyte-binding region which binds an analyte and causes the indicator to change conformation upon exposure to the analyte;

a donor fluorescent protein moiety covalently coupled to the binding protein moiety; and

an acceptor fluorescent protein moiety covalently coupled to the binding protein moiety,

wherein the donor moiety and the acceptor moiety change position relative to each other when the analyte binds to the analyte-binding region, altering fluorescence resonance energy transfer between the donor moiety and the acceptor moiety when the donor moiety is excited.

69. The indicator of claim 68, wherein the donor fluorescent protein moiety and the acceptor fluorescent protein moiety are *Aequorea*-related fluorescent protein moieties.

70. The indicator of claim 69, wherein the indicator further includes a target peptide moiety and a linker moiety that covalently couples the binding protein and the target peptide moiety, and the binding protein moiety further includes a peptide-binding region for binding the target peptide moiety.

71. The indicator of claim 70, wherein the indicator further comprises a localization sequence.

72. The indicator of claim 71, wherein the localization sequence is a nuclear localization sequence, an endoplasmic reticulum localization sequence, a peroxisome localization sequence, a mitochondrial localization sequence, or a localized protein.

WO 98/40477

PCT/US98/04978

- 81 -

73. The indicator of claim 70, wherein the donor fluorescent protein moiety is covalently coupled to the binding protein moiety and the acceptor fluorescent protein moiety is covalently coupled to the target peptide moiety.
74. The indicator of claim 70, wherein the linker moiety is a peptide moiety.
75. The indicator of claim 74, wherein the linker moiety includes between about 1 amino acid residue and about 30 amino acid residues.
76. The indicator of claim 74, wherein the indicator is a single polypeptide.
77. The indicator of claim 76, wherein one of the donor fluorescent protein moiety or the acceptor fluorescent protein moiety is covalently coupled to the carboxy terminus of the single polypeptide and the other of the donor fluorescent protein moiety or the acceptor fluorescent protein moiety is covalently coupled to the amino terminus of the single polypeptide.
78. The indicator of claim 69, wherein the binding protein moiety is calmodulin, a calmodulin-related protein moiety, cGMP-dependent protein kinase, a steroid hormone receptor, a ligand binding domain of a steroid hormone receptor, protein kinase C, inositol-1,4,5-triphosphate receptor, or recoverin.
79. The indicator of claim 70, wherein the binding protein moiety is calmodulin or a calmodulin-related protein moiety.
80. The indicator of claim 79, wherein the donor fluorescent protein moiety is P4-3, EBFP, or W1B, and the acceptor fluorescent protein moiety is S65T, EGFP, or 10c.

WO 98/40477

PCT/US98/04978

- 82 -

81. The indicator of claim 79, wherein the target peptide moiety is a calmodulin-binding domain of skMLCKp, smMLCK, CaMKII, Caldesmon, Calspermin, p-hosphofructokinase calcineurin, phosphorylase kinase, Ca²⁺-ATPase 59 kDa PDE, 60 kDa PDE, nitric oxide synthase, type I adenylyl cyclase, *Bordetella pertussis* adenylyl cyclase, Neuromodulin, Spectrin, MARCKS, F52, β -Adducin, HSP90a, HIV-1 gp160, BBMHBI, Dilute MHC, Mastoparan, Melittin, Glucagon, Secretin, VIP, GIP, or Model Peptide CBP2.
82. The indicator of claim 81, wherein the target peptide moiety is the calmodulin-binding domain of skMLCK.
83. The indicator of claim 82, wherein the linker moiety is -Gly-Gly-.
84. A fluorescent indicator comprising:
- a target peptide moiety;
 - a binding protein moiety having an analyte-binding region which binds an analyte and causes the indicator to change conformation upon exposure to the analyte, and a peptide-binding region for binding the target peptide moiety;
 - a linker moiety that covalently couples the binding protein and the target peptide moiety and is a peptide moiety;
 - a donor fluorescent protein moiety covalently coupled to the binding protein moiety; and
 - an acceptor fluorescent protein moiety covalently coupled to the target peptide moiety,
- wherein the donor moiety and the acceptor moiety change position relative to each other when the analyte binds to the analyte-binding region, altering fluorescence resonance energy transfer between the donor moiety and the acceptor moiety when the donor moiety is excited, and the indicator is a single polypeptide.

WO 98/40477

PCT/US98/04978

- 83 -

85. The indicator of claim 84, wherein the donor fluorescent protein moiety and the acceptor fluorescent protein moiety are *Aequorea*-related fluorescent protein moieties.
86. The indicator of claim 85, wherein the donor fluorescent protein moiety is P4-3, EBFP, or W1B, and the acceptor fluorescent protein moiety is S65T, EGFP, or 10c.
87. The indicator of claim 86, wherein the binding protein moiety is calmodulin or a calmodulin-related protein moiety.
88. The indicator of claim 87, wherein the target peptide moiety is M13.
89. The indicator of claim 88, wherein one of the donor fluorescent protein moiety or the acceptor fluorescent protein moiety is located at the carboxy terminus of the single polypeptide and the other of the donor fluorescent protein moiety or the acceptor fluorescent protein moiety is located at the amino terminus of the single polypeptide.
90. The indicator of claim 84, wherein the indicator further comprises a nuclear localization signal, an endoplasmic reticulum localization signal, a peroxisome localization signal, a mitochondrial localization signal, or a localized protein.

WO 98/40477

PCT/US98/04978

- 84 -

91. A method for determining the concentration of an analyte in a sample comprising:

contacting the sample with a fluorescent indicator comprising a binding protein moiety having an analyte-binding region which binds an analyte and causes the indicator to change conformation upon exposure to the analyte, a donor fluorescent protein moiety covalently coupled to the binding protein moiety, and an acceptor fluorescent protein moiety covalently coupled to the binding protein moiety, wherein the donor moiety and the acceptor moiety change position relative to each other when the analyte binds to the analyte-binding region, altering fluorescence resonance energy transfer between the donor moiety and the acceptor moiety when the donor moiety is excited; exciting the donor moiety; and

determining the degree of fluorescence resonance energy transfer in the sample corresponding to the concentration of the analyte in the sample.

92. The method of claim 91, wherein the step of determining the degree of fluorescence resonance energy transfer in the sample comprises measuring light emitted the acceptor fluorescent protein moiety.

93. The method of claim 91, wherein determining the degree of fluorescence resonance energy transfer in the sample comprises measuring light emitted from the donor fluorescent protein moiety, measuring light emitted from the acceptor fluorescent protein moiety, and calculating a ratio of the light emitted from the donor fluorescent protein moiety and the light emitted from the acceptor fluorescent protein moiety.

94. The method of claim 91, wherein the step of determining the degree of fluorescence resonance energy transfer in the sample comprises measuring the excited state lifetime of the donor moiety.

WO 98/40477

PCT/US98/04978

- 85 -

95. The method of claim 93, further comprising the steps of determining the concentration of the analyte at a first time after contacting the sample with the fluorescence indicator, determining the concentration of the analyte at a second time after contacting the sample with the fluorescence indicator, and calculating the difference in the concentration of the analyte at the first time and the second time, whereby the difference in the concentration of the analyte in the sample reflects a change in concentration of the analyte present in the sample.
96. The method of claim 95, further comprising the step of contacting the sample with a compound between the first time and the second time, whereby a difference in the concentration of the analyte in the sample between the first time and the second time indicates that the compound alters the presence of the analyte.
97. The method of claim 91, wherein the sample comprises an intact cell and the contacting step comprises incorporating the fluorescent indicator into the cell.
98. The method of claim 97, wherein step of incorporating the fluorescent indicator into the cell includes transfecting the cell with an expression vector comprising expression control sequences operably linked to a nucleic acid sequence coding for the expression of the fluorescent indicator.
99. The method of claim 91, wherein the analyte is calcium.
100. The method of claim 91, wherein the donor fluorescent protein moiety and the acceptor fluorescent protein moiety are *Aequorea*-related fluorescent protein moieties.

WO 98/40477

PCT/US98/04978


- 86 -

101. The method of claim 100, wherein the indicator further includes a target peptide moiety and a linker moiety that covalently couples the binding protein and the target peptide moiety and is a peptide moiety, and the binding protein moiety further includes a peptide-binding region for binding a target peptide moiety.
102. The method of claim 101, wherein the indicator is a single polypeptide.
103. The method of claim 91, wherein the binding protein moiety is calmodulin or a calmodulin-related protein moiety, the donor fluorescent protein moiety is P4-3, EBFP, or W1B, the acceptor fluorescent protein moiety is S65T, EGFP, or 10c, and the target peptide moiety is M13.
104. The method of claim 103, wherein the indicator further comprises a localization sequence.

WO 98/40477

PCT/US98/04978

1/17



151

WO 98/40477

PCT/US98/04978

2/17

FIG. 2a

50 a.a.

CaM-M13

GlyGly

HIS

P4-3 DONOR

XCaM

S65T ACCEPTOR

226 227

1 2

Ala Ala Arg Met His

GCT GCT CGC ATG CAT

20 21

1 2

Ile Ser Glu Leu Met Ser

ATC TCC GAG CTC ATG AGT

Sph1

Sac1

FIG. 2b

WO 98/40477

PCT/US98/04978

3/17




FIG. 3

WO 98/40477

PCT/US98/04978

4/17




FIG. 4a

WO 98/40477

PCT/US98/04978

5/17




FIG. 4b

WO 98/40477

PCT/US98/04978

6/17




FIG. 5

WO 98/40477

PCT/US98/04978

7/17

Fig. 6a





Fig. 6b



WO 98/40477

PCT/US98/04978

8/17

Fig. 6c





Fig. 6d

WO 98/40477

PCT/US98/04978

9/17



Fig. 6e




Fig. 6f

WO 98/40477

PCT/US98/04978

10/17

FIG. 7

(SEQ ID. NO:1)

ATGGTGAGCAAGGGCGAGGAGCTGTTACCGGGGTGGTGCCCATCCTGGTCCGAGCTGGACGGCGA
CGTAAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGA
CCCTGAAGTTCATCTGCACCACCGGCAAGCTGCCCGTGCCCTGGCCCAACCTCGTGACCAACCTG
ACCCATGGCGTGCAGTGCTTCAGCCGCTACCCCGACCACATGAAGCAGCAGCACTTCTTCAAGTC
CGCCATGCCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGA
CCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGAC
TTCAAGGAGGACGGCAACATCCTGGGGCACAAGCTGGAGTACAACCTTCAACAGCCACAACGTCTA
TATCATGGCCGACAAGCAGAAGAACGGCATCAAGGTGAACCTTCAAGATCCGCCACAACATCGAGG
ACGGCAGCGTGCAGCTCGCCGACCACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTG
CTGCCCCGACAACCACTACCTGAGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGA
TCACATGGTCCTGCTGGAGTTCGTGACCGCCGCCCGCATGCATGACCAACTGACAGAAGAGCAGA
TTGCAGAGTTCAAAGAAGCCTTCTCATTATTGACAAGGATGGGGACGGCACCATCACCACAAG
GAACTTGGCACCGTTATGAGGTGCTTGGACAAAACCCAAACGGAAGCAGAATTGCAGGATATGAT
CAATGAAGTCGATGCTGATGGCAATGGAACGATTTACTTTCCTGAATTTCTTACTATGATGGCTA
GAAAAATGAAGGACACAGACAGCGAAGAGGAAATCCGAGAAGCATTCGGTGTTTTTGACAAGGAT
GGGAACGGCTACATCAGCGCTGCTGAATTACGTCACGTCATGACCAACCTCGGGGAGAAGTTAAC
AGATGAAGGAGTTGATGAATGATAAGGGAAGCAGATATCGATGGTGATGGCCAAGTAAACTATG
AAGAGTTTGTACAAATGATGACAGCAAGGGGGGGAAGAGGCGCTGGAAGAAAACCTTCATTGCC
GTCAGCGCTGCCAACCGGTTCAAGAAGATCTCCGAGCTCATGCTGAGCAAGGGCGAGGAGCTGTT
CACCGGGGTGGTGCCCATCCTGGTCCGAGCTGGACGGCGACGTAAACGGCCACAAGTTCAGCGTGT
CCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACCGGC
AAGCTGCCCCGTGCCCTGGCCCAACCTCGTGACCAACCTGACCTACGGCGTGCACTGCTTCAGCCG
CTACCCCGACCACATGAAGCAGCAGCACTTCTTCAAGTCCGCCATGCCCCGAAGGCTACGTCCAGG
AGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGC
GACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGG
GCACAAGCTGGAGTACAACCAAGCCACAACGTCTATATCATGGCCGACAAGCAGAAGAAGC
GCATCAAGGTGAACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGCCGACCAC
TACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGCCCCGACAACCACTACCTGAGCAC
CCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTCCTGCTGGAGTTCGTGA
CCGCCGCCGGATCACTCTCGGCATGGACGAGCTGTACAAGTAA

WO 98/40477

PCT/US98/04978

11/17

FIG. 7 (continued)

(SEQ ID. NO:2)

MVSKGEELFTGVVPILVELDGDVNGHKFSVS GEGEGDATYGKLT LKFICTTGKLPVPWPTLVTTL
THGVQCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRA EVKFEGDTLVNRIELKGID
FKEDGNILGHKLEYNFN SHNVYIMADKQKNGIKVNFKIRHNI EDGSVQLADHYQQNTPIGDGPVL
LPDNHYLSTQSALSKDPNEKRDH MVLLFVTAARMHDQLTEEQIAEFKEAFSLFDKDG DGTITTK
ELGTVMRSLGQNPTEAELQDMINEVDADGNGTIYFPEFLTMMARKMKD TDSEEEIREAFRVFDKD
NGYISAAELRHVMTNLGEKLTDEEVD EMIREADIDGDGVNYEEFVQMMTAKGGKRRWKKNFIA
VSAANRFKKISELMVSKGEELFTGVVPILVELDGDVNGHKFSVS GEGEGDATYGKLT LKFICTTG
KLPVPWPTLVTTLT YGVQCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRA EVKFEG
DTLVNRIELKGIDFKEDGNILGHKLEYN FN SHNVYIMADKQKNGIKVNFKIRHNI EDGSVQLADH
YQQNTPIGDGPVLLPDNHYLSTQSALSKDPNEKRDH MVLLFVTAAGITLGMD ELYK*

WO 98/40477

PCT/US98/04978

12/17

FIG. 8

(SEQ ID. NO:3)

ATGGTGAGCAAGGGCGAGGAGCTGTTCAACGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGCGA
CGTAAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGA
CCCTGAAGTTCATCTGCACCACCGCAAGCTGCCCGTGCCCTGGCCACCCTCGTGACCACCCTG
ACCCATGGCGTGCACTGCTTCAGCCGCTACCCCGACCACATGAAGCAGCACGACTTCTTCAAGTC
CGCCATGCCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGA
CCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGAC
TTCAAGGAGGACGGCAACATCCTGGGGCACAAGCTGGAGTACAACCTTCAACAGCCACAACGTCTA
TATCATGGCCGACAAGCAGAAGAACGGCATCAAGGTGAACCTTCAAGATCCGCCACACATCGAGG
ACGGCAGCGTGCACTCGCCGACCACTACCAGCAGAACACCCCATCGGCGACGGCCCCGTGCTG
CTGCCCCGACAACCACTACCTGAGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGA
TCACATGGTCCTGCTGGAGTTCGTGACCGCCGCGCCGCATGCATGACCAACTGACAGAAGAGCAGA
TTGCAGAGTTCAAAGAAGCCTTCTCATTATTCGACAAGGATGGGGACGGCACCATCACCACAAAG
GAACTTGGCACCGTTATGAGGTGCTTGGACAAAACCCAACGGAAGCAGAAATTGCAGGATATGAT
CAATGAAGTCGATGCTGATGGCAATGGAACGATTTACTTTCTGAAATTTCTTACTATGATGGCTA
GAAAAATGAAGGACACAGACAGCGAAGAGGAAATCCGAGAAGCATTCGGTGTTTTTGACAAGGAT
GGGAACGGCTACATCAGCGCTGCTGAATTACGTACGTCATGACAAACCTCGGGGAGAAGTTAAC
AGATGAAGAAGTTGATGAAATGATAAGGGAAGCAGATATCGATGGTGATGGCCAAGTAAACTATG
AAGAGTTTGTACAAATGATGACAGCAAAGGGGGGAAGAGGCGCTGGAAGAAAACTTCATTGCC
GTCAGCGCTGCCAACCAGTTCAAGAAGATCTCCGAGCTCATGGTGACAAAGGGCGAGGAGCTGTT
CACCGGGGTGGTGCCCATCCTGGTCCAGCTGGACGGCGACGTAAACGGCCACAAGTTCAGCGTGT
CCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACCGGC
AAGCTGCCCCGTGCCCTGGCCACCCTCGTGACCACCCTGACCTACGGCGTGCACTGCTTCAGCCG
CTACCCCGACCACATGAAGCAGCACGACTTCTTCAAGTCCGCCATGCCCGAAGGCTACGTCCAGG
AGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGC
GACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGG
GCACAAGCTGGAGTACAACCTACAACAGCCACAACGTCTATATCATGGCCGACAAGCAGAAGAAG
GCATCAAGGTGAACCTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCACTCGCCGATCCAC
TACCAGCAGAACACCCCATCGGCGACGGCCCCGTGCTGCTGCCCCGACAACCACTACCTGAGCAC
CCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTCCTGCTGGAGTTCGTGA
CCGCCGCGGGATCACTCTCGGCATGGACGAGCTGTACAAGCCAAAAAGAAGAGAAGGTGGAA
GACGCTTAA

WO 98/40477

PCT/US98/04978

13/17

FIG. 8 (continued)

(SEQ ID. NO:4)

MVSKGEELFTGVVPILVELDGDVNGHKFSVS GEGEGDATYGKLT LKFICTTGKLPVPWPTLVTTL
THGVQCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTLVNRIELKGID
FKEDGNILGHKLEYNFSHNVIYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVL
LPDNHYLSTQSALS KDPNEKRDHMLLEFVTAARMHDQLTEEQIAEFKEAFLFDKDGDTITTK
ELGTVMRSLGQNPTEAELQDMINEVDADGNGTIYFPEFLTMARKMKD TDSEEEIREAFRVFDKD
GNGYISAAELRHVMTNLGEKLTDEEVDEMIREADIDGDCQVNYEEFVQMMTAKGGKRRWKKNFIA
VSAANRFKKISELMVSKGEELFTGVVPILVELDGDVNGHKFSVS GEGEGDATYGKLT LKFICTTG
KLPVPWPTLVTTLT YGVQCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFEG
DTLVNRIELKGIDFKEDGNILGHKLEYNFSHNVIYIMADKQKNGIKVNFKIRHNIEDGSVQLADH
YQQNTPIGDGPVLLPDNHYLSTQSALS KDPNEKRDHMLLEFVTAAGITLGMD ELYKPKKKRKVE
DA*

WO 98/40477

PCT/US98/04978

14/17

FIG. 9

(SEQ ID. NO:5)

ATGGTGAGCAAGGGCGAGGAGCTGTTACCCGGGGTGGTGCCCATCCTGGTCCGAGCTGGACGGCGA
CGTAAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGA
CCCTGAAGTTCATCTGCACCACCGGCAAGCTGCCCCGTGCCCTGGCCCCACCCTCGTGACCACCCTG
ACCCATGGCGTGCAGTGCTTCAGCCGCTACCCCGACCACATGAAGCAGCAGCACTTCTTCAAGTC
CGCCATGCCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGA
CCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGAC
TTCAAGGAGGACGGCAACATCCTGGGGCACAAGCTGGAGTACAACCTTCAACAGCCACAACGTCTA
TATCATGGCCGACAAGCAGAAGAACGGCATCAAGGTGAACCTTCAAGATCCGCCACAACATCGAGG
ACGGCAGCGTGACGCTCGCCGACCACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTG
CTGCCCCGACAACCACTACCTGAGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGA
TCACATGGTCTGCTGGAGTTCGTGACCGCCGCCCGCATGCATGACCAACTGACAGAAGAGCAGA
TTGCAGAGTTCAAAGAAGCCTTCTCATTATTCGACAAGGATGGGGACGGCACCATCACCACAAAG
GAACTTGGCACCGTTATGAGGTCGCTTGGACAAAACCCAACGGAAGCAGAATTGCAGGATATGAT
CAATGAAGTCGATGCTGATGGCAATGGAACGATTTACTTTCTGAATTTCTTACTATGATGGCTA
GAAAAATGAAGGACACAGACAGCGAAGAGGAAATCCGAGAAGCATTCCGTGTTTTTTGACAAGGAT
GGGAACGGCTACATCAGCGCTGCTCAGTTACGTACGTCATGACAAAACCTCGGGGAGAAGTTAAC
AGATGAAGAAGTTGATGAATGATAAGGGAAGCAGATATCGATGGTGATGGCCAAGTAACTATG
AAGAGTTTGTACAAATGATGACAGCAAAGGGGGGGAAGAGGCGCTGGAAGAAAACCTTCATTGCC
GTCAGCGCTGCCAACCGGTTCAAGAAGATCTCCGAGCTCATGGTGAGCAAGGGCGAGGAGCTGTT
CACCGGGGTGGTGCCCATCCTGGTCCGAGCTGGACGGCGACGTAAACGGCCACAAGTTCAGCGTGT
CCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACCGGC
AAGCTGCCCCGTGCCCTGGCCCACCCTCGTGACCACCCTGACCTACGGCGTGACGTGCTTCAGCCG
CTACCCCGACCACATGAAGCAGCAGCACTTCTTCAAGTCCGCCATGCCCGAAGGCTACGTCCAGG
AGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGC
GACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGG
GCACAAGCTGGAGTACAACCAACAGCCACAACGTCTATATCATGCGCGACAAGCAGAAGAACG
GCATCAAGGTGAACCTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGACGCTCGCCGACCAC
TACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGCCCCGACAACCACTACCTGAGCAC
CCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTCTCTGCTGAGTTCGTGA
CCGCCGCCGGGATCACTCTCGGCATGGACGAGCTGTACAAGTAA

WO 98/40477

PCT/US98/04978

15/17

FIG. 9 (continued)

(SEQ ID. NO:6)

MVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTCLKFICTTGKLPVPWPTLVTTL
THGVQCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTLVNRIELKGID
FKEDGNILGHKLEYNFNHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVL
LPDNHYLSTQSALS KDPNEKRDHMLLEFVTAARMHDQLTEEQIAEFKEAFSLFDKDGDTITTK
ELGTVMRSLGQNPTEAELQDMINEVDADGNGTIYFPEFLTMMARKMKDSEEEIREAFRVFDKD
GNGYISAAQLRHVMTNLGEKLTDEEVDEMIREADIDGDGQVNYEEFVQMMTAKGGKRRWKKNFIA
VSAANRFKKISELMVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTCLKFICTTG
KLPVPWPTLVTTLTYGVCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFEG
DTLVNRIELKGIDFKEDGNILGHKLEYNHNHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADH
YQQNTPIGDGPVLLPDNHYLSTQSALS KDPNEKRDHMLLEFVTAAGITLGMDELYK*

WO 98/40477

PCT/US98/04978

16/17

FIG. 10

(SEQ ID. NO:7)

ATGCTGCTGCCCCGTCCCCCTGCTGCTGGGCCTGCTGGGCGCCGCCGCCGACGTGAGCAAGGGCGA
GGAGCTGTTCAACGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGCGACGTAAACGGCCACAAGT
TCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGC
ACCACCGGCAAGCTGCCCCGTGCCCTGGCCACCCCTCGTGACCACCCTGACCCATGGCGTGCAGTG
CTTCAGCCGCTACCCCGACCACATGAAGCAGCAGACTTCTTCAAGTCCGCCATGCCCCGAAGGCT
ACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGACCCGCGCCCGAGGTGAAG
TTCGAGGGCGACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAA
CATCCTGGGGCACAAGCTGGAGTACAACCTTCAACAGCCACAACGTCTATATCATGGCCGACAAGC
AGAAGAACGGCATCAAGGTGAACCTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTC
GCCGACCACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGCCCGACAACCACTA
CCTGAGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTCCTGCTGG
AGTTCGTGACCGCCGCCCGCATGCATGACCAACTGACAGAAGAGCAGATTGCAGAGTTCAAAGAA
GCCTTCTCATTATTCGACAAGGATGGGGACGGCACCATCACCACAAGGAACTTGGCACCGTTAT
GAGGTGCTTGGACAAAACCCAACGGAAGCAGAATTGCAGGATATGATCAATGAAGTCGATGCTG
AEGGCAATGGAACGATTTACTTTCCTGAATTTCTTACTATGATGGCTAGAAAATGAAGGACACA
GACAGCGAAGAGGAAATCCGAGAAGCATTCCGTGTTTTTGACAAGGATGGGAACGGCTACATCAG
CGCTGCTCAGTTACGTACGTCATGACAAACCTCGGGGAGAAGTTAACAGATGAAGAAGTTGATG
AAATGATAAGGGAAGCAGATATCGATGGTGATGGCCAAGTAACTATGAAGAGTTTGTAACAATG
ATGACAGCAAAGGGGGGGAAGAGGCGCTGGAAGAAAACTTCATTGCCGTGAGCGCTGCCAACCG
GTTCAAGAAGATCTCCGAGCTCATGGTGAGCAAGGGCGAGGAGCTGTTACCGGGGTGGTGCCCA
TCCTGGTTCGAGCTGGACGGCGACGTAAACGGCCACAAGTTCAGCGTGTCGGCGAGGGCGAGGGC
GATGCCA.CCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACCGCAAGCTGCCCCGTGCCCTG
GCCACCCCTCGTGACCACCCTGACCTACGGCGTGCAAGTCTCAGCCGCTACCCCGACCACATGA
AGCAGCACGACTTCTTCAAGTCCGCCATGCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTC
AAGGACGACGGCAACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCG
CATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGGGCACAAGCTGGAGTACA
ACTACAACAGCCACAACGTCTATATCATGGCCGACAAGCAGAAGAACGGCATCAAGGTGAACCTTC
AAGATCCGCCACAACATCGAGGACGGCAGCGTGCAAGCTCGCCGACCACTACCAGCAGAACACCCC
CATCGGCGACGGCCCCGTGCTGCTGCCCCACAACCACTACCTGAGCACCCAGTCCGCCCTGAGCA
AAGACCCCAACGAGAAGCGCGATCACATGGTCCTGCTGGAGTTCGTGACCGCCGCGGGATCACT
CTCGGCAAGGACGAGCTGTAA

WO 98/40477

PCT/US98/04978

17/17

FIG. 10 (continued)

(SEQ ID. NO:8)

MLLPVPLLLGLLGAAADVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTCLKFIC
TTGKLPVPWPTLVTTLTGTVQCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVK
FEGDTLVNRIELKGIDFKEDGNILGHKLEYNFNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQL
ADHYQONTPIGDGPVLLPDNHYLSTQSALSKDPNEKRDHMLLEFVTAARMHDQLTEEQIAEFKE
AFSLFDKDGDTITTKELGTVMRSLGQNPTEAELQDMINEVDADGNGTIYFPEFLTMMARKMKDT
DSEEEIREAFRVFDKDGNGYISAAQLRHVMTNLGEKLTDEEVDEMIREADIDGDGQVNYEEFVQM
MTAKGGKRRWKKNFIAVSAANRFKKISELMVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEG
DATYGKLTCLKFICTTGKLPVPWPTLVTTLTGTVQCFSRYPDHMKQHDFFKSAMPEGYVQERTIFF
KDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNFNSHNVYIMADKQKNGIKVNF
KIRHNIEDGSVQLADHYQONTPIGDGPVLLPDNHYLSTQSALSKDPNEKRDHMLLEFVTAAGIT
LGKDEL*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/04978

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 15/00, 15/09; C07H 21/02, 21/04

US CL : 435/320.1, 325; 536/22.1, 23.1, 23.4

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/320.1, 325; 536/22.1, 23.1, 23.4

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

None

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CAS ONLINE

Search terms: nucleic acid, expression vector, host cell line, fluorescent protein, energy transfer

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4,868,103 A (STAVRIANOPOULOS et al.) 19 September 1989, column 5, line 20 to column 7, line 35; column 12, line 40 to column 17, line 58; column 23, line 6 to column 28, line 25.	1-21, 23, 25-33, 51, 53-61
A	US 5,134,232 A (TSIEN et al.) 28 July 1992, see entire document.	1-21, 23, 25-33, 51, 53-61
Y	US 5,439,797 A (TSIEN et al) 08 August 1995, column 2, line 36 to column 8, line 31; column 10, line 26 to column 11, line 68.	1-21, 23, 25-33, 51, 53-61

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

11 MAY 1998

Date of mailing of the international search report

15 JUL 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

P. ACHUTAMURTHY

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/04978

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-21, 23, 25-33, 40-51, 53-61

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/04978

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1.

Group I, claim(s) 1-21, 23, 25-33, 40-51, and 53-61, drawn to nucleic acid, an expression vector comprising the nucleic acid, and a host cell line comprising the vector.

Group II, claim(s) 24, 34-39, 52, and 62-67, drawn to a transgenic animal.

Group III, claim(s) 22, drawn to a system for monitoring protei-protein association.

Group IV, claims 68-90, drawn to a fluorescent indicator.

Group V, claims 91-104, drawn to a method of analyte determination using a fluorescent indicator.

The inventions listed as Groups I-V do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: Claims of Groups I-V are not related because they are drawn to different products and methods.

The product disclosed in Group II, is a transgenic animal which is distinct from an isolated cell line carrying an expression vector comprising a nucleic acid. Group III is drawn to a system comprising two fluorescent protein moieties and does not require the elements of either group I or Group II. It may comprise protein moieties which are structurally and functionally different from the protein expressed by the vector recited in group I. Similarly group IV is drawn to fluorescent protein which may be different from the protein expressed by the vector of group I. The method of group V can be practiced with a combination of protein moieties which are materially, structurally, and functionally different from the protein required in groups I-IV.